

## ALGEBRA – MODEL NO

1

**[Q1] A) Choose the correct answer:**

(1) The two straight lines which represent the two equations:

$X = 3, Y = 5$  are .....

- a) Perpendicular                      c) Coincide  
b) Parallel                              d) Intersect and not perpendicular

(2) The equation  $\frac{1}{x} + \frac{1}{y} = 3$  of ..... Degree ( $X \neq Y \neq 0$ )

- a) First                      b) Second                      c) Third                      d) Fourth

(3) Number of solution of the equation  $2X - 6 = 0$  in  $\mathcal{R}^2$  is.....

- a) 1                      b) 2                      c) 3                      d) Infinite

[B] By using a general formula, find in  $\mathcal{R}$  the solution set of the equation  $X^2 - 2X - 6 = 0$ , approximating the result to nearest two decimal places.

**[Q2] Choose the correct answer:**(1) A number formed from two digits, its units digit = its tens digit =  $X$ , then the number is .....

- a)  $X^2$                       b)  $2X$                       c)  $11X$                       d)  $10X^2$

(2) If  $n(x) = \frac{x-3}{x+2}$ ,  $n^{-1}(K) = \frac{7}{2}$ , then  $K = \dots\dots\dots$ ,  $X \notin \{3, -2\}$ 

- a)  $-4$                       b)  $5$                       c)  $-5$                       d)  $-\frac{8}{9}$

(3) If  $A, B$  are two mutually exclusive events from the sample space of a random experiment, then  $A \cap B = \dots\dots\dots$ 

- a)  $\emptyset$                       b)  $S$                       c) Zero                      d) 1

[B] Find  $n(x)$  in the simplest form and showing its domain:

$$n(x) = \frac{x^2 - 2x - 15}{x^2 - 9} \div \frac{2x - 10}{x^2 - 6x + 9}$$

[Q3]

[A] If the set of zeroes of  $\mathcal{F}: \mathcal{F}(x) = aX^2 + bx + 15$  is  $\{3, 5\}$ . Find the value of each of  $a, b$

---

[B] If  $n_1(x) = \frac{x^2-4}{x^2+x-6}, n_2(x) = \frac{x^2-x-6}{x^2-9}$

Show that if  $n_1(x) = n_2(x)$  or not?

Find the common domain in which that  $n_1(x) = n_2(x)$ ?

---

[Q4]

[A] Find in the simplest form and showing its domain:

$$n(x) = \frac{x^2+3x+9}{x^3-27} + \frac{(x-4)^2}{x^2-7x+12}$$


---

[B] A right angled-triangle, the length of one of right angled sides is 5 cm, and its perimeter 30 cm, find its surface area?

---

[Q5]

[A] If  $A, B$  are two events of the sample space of a random experiment, and  $P(A) = 0.6, P(B) = 0.7, P(A \cap B) = 0.4$ , Find:

①  $P(A - B)$

② The probability of the occurrence of one of the two events at least

---

[B] IF  $\frac{k+5-x^2}{x^2-3x}$  is additive inverse of the fraction  $\frac{x}{x-3}$ .

Find the value of  $K$ .

---

*End of the questions*



## ALGEBRA – MODEL NO 2

[Q1]: A) Choose the correct answer:

(1) The equation  $3X + 4Y + XY = 5$  of ..... degree

- a) First                      b) Second                      c) Third                      d) Fourth

(2) The two straight lines which represented two equations  $3X + 5Y = 0$ ,  $5X - 3Y = 0$  intersect at the point .....

- a)  $(0, 0)$                       b)  $(-5, 3)$                       c)  $(3, 5)$                       d)  $(-3, -5)$

(3) If  $n(x) = \frac{x-2}{x+1}$ , then  $n^{-1}(2)$  .....

- a) = zero                      b) = 2                      c) = 3                      d) undefined

B): By using a general formula, find in  $\mathcal{R}$  the solution set of the equation  $X(X - 1) = 4$ , approximating the result to one decimal place.

[Q2]: A) Choose the correct answer:

(1) If  $XY = 3$ ,  $XY^2 = 12$ , then  $Y =$  .....

- a) 4                      b) 2                      c) -2                      d)  $\pm 2$

(2) If A, B are two mutually exclusive events, then  $P(A \cap B) =$  ...

- a)  $\emptyset$                       b) 1                      c) 0.5                      d) Zero

(3) The domain of  $\mathcal{F}$ :  $\mathcal{F}(x) = X^2 + 4$  is .....

- a)  $\mathcal{R} - \{2, -2\}$                       b)  $\{2, -2\}$                       c)  $\mathcal{R}$                       d)  $\emptyset$

B): Prove that  $n_1(x) = n_2(x)$  where:

$$n_1(x) = \frac{2x}{2x+8}, \quad n_2(x) = \frac{x^2+4x}{x^2+8x+16}$$

[Q3]

A): If the domain of  $n(X) = \frac{b}{x} + \frac{9}{x+a}$  is  $\mathbb{R} - \{0, 4\}$ ,  $n(5) = 2$ .

Find the value of A , B

---

B): Two acute angles in right angle triangle, the difference between them is  $50^\circ$ , find the measure of each angle.

---

[Q4]

A): Put in the simplest form and showing its domain:

$$n(X) = \frac{x^2 - 2x}{x^2 - 3x + 2} - \frac{4 - x^2}{x^2 + x - 2}$$


---

B): Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of two equations:

$$Y + 2X = 7 \quad , \quad (Y + 2X - 8)^2 + X^2 = 5$$


---

[Q5]

A): Put in the simplest form and showing its domain:

$$n(X) = \frac{x^3 - 8}{x^2 + x - 6} \times \frac{x + 3}{x^2 + 2x + 4}$$


---

B): If A , B are two events of the sample space of a random experiment, and  $P(A) = 0.5$ ,  $P(B) = 0.4$ ,  $P(A \cap B) = 0.1$ , Find:

①  $P(A \cup B)$

②  $P(A - B)$

◆ ◆ ◆

*End of the questions*



## ALGEBRA – MODEL No

3

[Q1] A) Choose the correct answer:

(1) The solution set of two equations  $X - 3 = 0$ ,  $Y = 4$  in  $\mathcal{R} \times \mathcal{R}$  is ...

- a)
- $\{3, 4\}$
- b)
- $\{(3, 4)\}$
- c)
- $\{(4, 3)\}$
- d)
- $\emptyset$

(2) If  $A, B$  are two events in sample space of a random experiment,  $A \subset B$ , then  $P(A \cup B) = \dots\dots\dots$ 

- a)
- $P(B)$
- b)
- $P(A)$
- c)
- $P(A \cap B)$
- d) Zero

(3) If  $3^Y \times 5^Y = 225$ , then  $Y = \dots\dots\dots$ 

- a) 2      b) 15      c) Zero      d) 20

B):Find in  $\mathcal{R} \times \mathcal{R}$  the solution set of two equations:

$$3X - Y = 5, X + 2Y = 4$$

[Q2] A) Choose the correct answer:

(1) The domain of the additive inverse of  $n(x) = \frac{x+2}{x-3}$ 

- a)
- $\mathcal{R} - \{3\}$
- b)
- $\mathcal{R} - \{-2\}$
- c)
- $\mathcal{R} - \{-2, 3\}$
- d)
- $\mathcal{R}$

(2) The set of zeroes of  $\mathcal{F}(x) = X^2 + 9$  in  $\mathcal{R}$  is .....

- a)
- $\mathcal{R}$
- b)
- $\emptyset$
- c)
- $\{3\}$
- d)
- $\{3, -3\}$

(3) The curve  $Y = aX^2 + bx + C$  cut  $Y$ -axis at the point .....

- a)
- $(0, b)$
- b)
- $(b, 0)$
- c)
- $(c, 0)$
- d)
- $(0, 7)$

B):

Put in the simplest form:  $n(X) = \frac{x^2+x}{x^2-1} - \frac{5-x}{x^2-x+5}$

And showing its domain

[Q3]

A) If  $A$ ,  $B$  are two events of the sample space of a random experiment, and  $P(A) = 0.6$ ,  $P(B) = 0.5$ ,  $P(A \cap B) = 0.3$ , Find

①  $P(A \cup B)$

②  $P(B^c)$

B) Put in the simplest form:  $n(X) = \frac{x^3 - 1}{x^2 - 2x + 1} - \frac{2x - 2}{x^2 + x + 1}$

And showing its domain

[Q4]

A) Prove that  $n_1(x) = n_2$

Where  $n_1(x) = \frac{x^2 - x}{x^3 - 2x^2}$ ,  $n_2(x) = \frac{x^2 - 3x + 2}{x^3 - 4x^2 + 4x}$

B) By using a general formula, find in  $\mathcal{R}$  the solution set of the equation  $2X^2 + 4X + 1 = 0$ , approximating the result into two decimal places.

[Q5]

A) Find the solution set of two equations in  $\mathcal{R} \times \mathcal{R}$

$$X - Y = 0 \quad , \quad X = \frac{4}{y}$$

B) If  $n(X) = \frac{x^2 - 2x}{(x-2)(x^2+2)}$  Find:

①  $n^{-1}(x)$  showing its domain

② If  $n^{-1}(x) = 3$ , find the value of  $X$ .

◆◆◆

End of the questions



## ALGEBRA — MODEL No 4

[Q1] A) Choose the correct answer:

(1) If  $S$  is sample space, then  $P(S) = \dots\dots\dots$

- a) 1                      b) 0                      c) Half                      d) -1

(2) If the algebraic fraction  $\frac{x-a}{x+3}$  is additive inverse to  $\frac{x+3}{x+5}$  then  $a = \dots\dots\dots$

- a) -5                      b) -3                      c) 5                      d) 3

(3) If  $x^2 + y^2 = 5xy$ , then  $\frac{x^2}{y^2} + \frac{y^2}{x^2} = \dots\dots\dots$

- a) 32                      b) 23                      c) -32                      d) -23

B): By using a general formula, find in  $\mathcal{R}$  the solution set of the equation  $\frac{2}{x^2} = 1 - \frac{2}{x}$ , approximating the result to nearest two decimal places.

[Q2] A) Choose the correct answer:

(1) If the two equation :  $3x - 5y = 8$ ,  $2x + ky = m$  has infinite solution in  $\mathcal{R} \times \mathcal{R}$  then,  $9km = \dots\dots\dots$

- a)  $-\frac{10}{3}$                       b)  $\frac{16}{3}$                       c) -16                      d) -160

(2) If the set of zeros of the function  $f$  where  $F(x) = kx + 3$  is  $\emptyset$  then  $k = \dots\dots\dots$

- a) -3                      b) 3                      c) 0                      d) 1

(3) The function  $f : f(x) = \frac{x-2}{x-5}$  has additive inverse in the domain

- a)  $\mathcal{R} - \{2\}$                       b)  $\mathcal{R} - \{5\}$                       c)  $\mathcal{R} - \{5, -2\}$                       d)  $\mathcal{R} - \{5, 2\}$

B): If:  $n_1(x) = \frac{3x-6}{x^2-4}$ ,  $n_2(x) = \frac{3x+3}{x^2+3x+2}$  prove that  $n_1(x) = n_2(x)$   
For all values of  $x$  in common domain and Find this domain?

[Q3]

A) A rectangle the length of its diagonal 5 cm, its perimeter 14 cm.  
find its dimensions.

B) Put in the simplest form and showing the domain:

$$n(x) = \frac{x-6}{2x^2-15x+18} - \frac{x-5}{15-13x+x^2}$$

[Q4] If the set of zeros of the function  $n$  where  $N(x) = \frac{x^2-ax+9}{bx+4}$   
is  $\{3\}$ , its domain  $R - \{2\}$  find the value of  $a$  and  $b$ ?

B) If  $A, B$  are two events of the sample space of a random experiment, and  $P(A) = P(A^c)$ ,  $P(B) = \frac{1}{3}$ ,  $P(A-B) = \frac{5}{12}$  Find:

- ① The probability of non-occurrence of  $A, B$  together.
- ② The probability of occurrence at least one of them.


[Q5] A) Find in the simplest form:  $n(X) = \frac{x^2-2x-15}{x^2-9} \div \frac{x^2-25}{x^2-3x}$   
And showing its domain. If  $n(k) = \frac{1}{3}$ , find the value of  $K$

B) In the opposite figure:

The curve of  $F: F(x) = kx^2 + mx + n$

Cut  $Y$ -axis in  $c(0,6)$ ,  $A(1,4)$  is the vertex curve

Find the value of  $K, m, n$



End of the question



## ALGEBRA – MODEL No

5

**[Q1] A) Choose the correct answer:**

(1) If A is an event in a sample space of a random experiment, and  $P(A \cup A^c) = \dots\dots\dots$

- a) 1                      b) 0                      c)  $\frac{1}{2}$                       d) -1

(2) The set of zeros of  $\mathcal{F}(x) = \frac{x^2 - x - 2}{x^2 - 4}$  is  $\dots\dots\dots$

- a)  $\{-1, 2\}$               b)  $\{-2, 2\}$               c)  $\{-1\}$                       d)  $\{-2\}$

(3) Two straight lines  $3X + 5Y = 0$ ,  $5X - 3Y = 0$  intersect in  $\dots\dots\dots$

- a) Origin point      b) First quadrant              c) Second quadrant              d) Fourth quadrant

**B):** By using a general formula and, find in  $\mathcal{R}$  the solution set of the equation  $X + \frac{4}{x} = 6$ , approximating the result to nearest three decimal places

**[Q2] A) Choose the correct answer:**

(1) In the equation:  $aX^2 + 6X + 3 = 0$ , has no solution in  $\mathcal{R}$  then  $a \in \dots\dots\dots$

- a)  $] -\infty, 3[$               b)  $] 3, \infty [$               c)  $\{3\}$                       d)  $\{-3, 3\}$

(2) If  $-1 + x^2 = 3x$ , then  $x + \frac{1}{x} = \dots\dots\dots$

- a) 1                      b) 3                      c) -1                      d) -3

(3) If  $\mathcal{F}(x) = \frac{x^2 - x}{x^2 - 1}$ ,  $\mathcal{F}^{-1}(k) = 3$ , then  $K = \dots\dots\dots$

- a)  $-\frac{3}{2}$                       b)  $\frac{1}{2}$                       c)  $\frac{3}{4}$                       d)  $\frac{4}{3}$

B) A rhombus the difference between lengths of its diagonals 4 cm, its perimeter 40 cm. find its diagonal lengths.

---

[Q3]

A) Find in the simplest form:  $n(x) = \frac{2x+6}{x^2+x-6} + \frac{3x-4}{2x^2-5x+6}$   
And showing its domain

---

B) If  $n_1(x) = \frac{x^2-4}{x^2+x-6}$ ,  $n_2(x) = \frac{x^2-x-6}{x^2-9}$  Prove that  $n_1(x) = n_2(x)$  ?

---

[Q4] A) Find in the simplest form:  $n(x) = \frac{x^2-2x-15}{x^2-9} \div \frac{2x-10}{x^2-6x+9}$   
And showing its domain

---

B) If the domain of  $n(x) = \frac{a}{x-3} + \frac{4}{x+b}$  is  $\mathcal{R} - \{3, -4\}$ ,  $n(2) = 7$ , find the value of a, b ?

---

[Q5] A) If A,B are two events of the sample space of a random experiment, and  $P(A) = \frac{1}{2}$ ,  $P(B) = \frac{2}{5}$ ,  $P(A \cap B) = \frac{1}{10}$  Find:

①  $P(A \cup B)$

②  $P(B - A)$

---


B) In the opposite figure:

If the equation of  $\overleftrightarrow{AB}$ :  $X + Y = 3$ ,

Equation of  $\overleftrightarrow{CD}$ :  $2X + Y + 4 = 0$

Find the area of the shaded part

*End of the question*







# Prep. 3 - Model (01)



## [Q1] A) Choose the correct answer:

(1) If the domain of  $n(x) = \frac{x-1}{x-a}$  is  $R - \{2\}$ , then  $a = \dots\dots\dots$

- a) -2                      b) -1                      c) 1                      d) 2

(2) If  $X - Y = 1$ ,  $(X - Y)^2 + Y = 1$ , then  $X = \dots\dots\dots$

- a) -2                      b) -1                      c) 1                      d) 2

(3) If  $A$  is an event in a sample space of a random experiment, and  $P(A) = 4 P(A^c)$ , then  $P(A) = \dots\dots\dots$

- a) 4                      b) 1                      c)  $\frac{4}{5}$                       d)  $\frac{1}{4}$

**B):** By using a general formula and without using calculator, find in  $R$  the solution set of the equation  $X^2 - 8X + 3 = 0$ , and  $\sqrt{13} \simeq 3.6$

## [Q2] A) Choose the correct answer:

(1) The two equations  $3X - 2Y = 5$ ,  $3X - 2Y = K$ , have infinite number of solutions when  $k = \dots\dots\dots$

- a) 3                      b) 2                      c) -5                      d) 5

(2) If  $X = 1$  is one of the set of zeroes of  $F(X) = X^2 - 3X + C$ , then  $C = \dots\dots\dots$

- a) Zero                      b) 1                      c) 2                      d) 3

(3) Which of the following algebraic fractional in the simplest form?

- a)  $\frac{x+1}{x^2+1}$                       b)  $\frac{x+1}{x^2-1}$                       c)  $\frac{x}{x^2}$                       d)  $\frac{x}{x^2+x}$

**B):** Find each of  $n_1(x) = \frac{2x}{2x+4}$ ,  $n_2(x) = \frac{x^2+2x}{x^2+4x+4}$  in the simplest form

showing the domain of each one then show that if  $n_1 = n_2$  or not?

Give the reason

**[Q3]****A)** Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of two equations:

$$X = 2Y + 3, Y^2 - X = 0$$


---

**B)** Find in the simplest form and showing its domain:

$$n(x) = \frac{x^2 - 9}{x^2 - x - 6} - \frac{x^2 - 4x}{x^2 - 2x - 8}$$


---

**[Q4]****A)** If  $n$  is an algebraic fractional function where  $n(x) = \frac{2+b}{x+4}$ , find the domain of  $n$ . If  $n(5) = 1$  **find** the value of expression:  $2b - 11$ **B)** A bag contains **20** cards numbered from **1** to **20**, on card is chosen randomly, **find** the probability of chosen card has a number:

① Divisible by 3

② Odd and divisible by 5

**[Q5]****A)** Find algebraically the solution set of two equations:

$$X + Y - 5 = 0, 3X + Y = 17$$


---

**B)** Find in the simplest form:  $n(X) = \frac{x^2 - 3x + 2}{x^2 + x - 6} \times \frac{x^2 + 2x}{x^2 + x - 2}$ And show its domain, and then find if possible the value of  $n(1)$ 

\*\*\* End of the questions \*\*\*





# Prep. 3 - Model (02)



## [Q1] A) Choose the correct answer:

(1) The set of zeroes of  $F(x) = X + 3$  is .....

- a)  $\mathbb{R}$                       b)  $\mathbb{R} - \{3\}$                       c)  $\{3\}$                       d) 3

(2) The two straight line  $X = 4$ ,  $Y = 3$  intersecting at point ....

- a)  $(4, 3)$                       b)  $(0, 0)$                       c)  $(3, 4)$                       d)  $(-3, -4)$

(3) If  $X$ ,  $Y$  are two mutually exclusive events in the sample space of a random experiment, then  $P(X \cap Y) = \dots\dots\dots$

- a)  $\emptyset$                       b) Zero                      c)  $\{\}$                       d) 1

**B):** find the solution set of two equations:

$$X - Y = 0 \quad , \quad XY = 4$$

## [Q2] A) Choose the correct answer:

(1) The two first degree equations in one variable which have infinite solution are represented graphically with two straight lines are.....

- a) Parallel                      b) Interest at one point                      c) Congruent                      d) Disjoint

(2) If  $F(x) = \frac{7+X}{7-X}$ , where  $x \in \mathbb{R} - \{7, -7\}$ , then  $F(-2) = \dots\dots\dots$

- a)  $\frac{-1}{f(-2)}$                       b)  $\frac{-1}{f(2)}$                       c)  $\frac{1}{f(2)}$                       d)  $\frac{1}{f(-2)}$

(3) If the domain of function  $n$  where  $n(x) = \frac{x-2}{x^2+K}$  is  $\mathbb{R}$

Then  $K \dots\dots\dots$  zero

- a)  $=$                       b)  $<$                       c)  $>$                       d)  $\leq$

**B):** A rectangle with a length is more than their widths by 5 cm. if the perimeter of the rectangle is 18 cm, find the area of rectangle

**[Q3]****A)** Find  $n(x)$  in the simplest form showing its domain:

$$n(x) = \frac{x^2 - 3x + 2}{x^2 - 1} \times \frac{3x - 15}{x^2 - 4x - 5}$$


---

**B)** By using a general formula and without using calculator, find in  $\mathbb{R}$  the solution set of the equation  $X + \frac{1}{X} = 5$ , approximating the result to two decimal places. and  $\sqrt{17} \simeq 4.12$ **[Q4]****A)** Find  $n(x)$  in the simplest form showing its domain:

$$n(x) = \frac{x^2 + x + 1}{x^3 - 1} \div \frac{x^2 - x}{x^2 - 2x + 1}$$


---

**B)** Find in the simplest form:  $n(X) = \frac{x^2 - 9}{x^2 - x - 6} + \frac{4x - x^2}{x^2 - 2x - 8}$   
And show its domain, and then find if possible the value of  $n(3)$ **[Q5]****A)** If  $A, B$  are two events of the sample space of a random experiment,

$$\text{and } P(A) = \frac{1}{2}, P(B) = \frac{2}{5}, P(A \cap B) = \frac{1}{10}, \text{ Find}$$

①  $P(A \cup B)$

②  $P(B - A)$

**B)** If  $n_1, n_2$  are two functions, Prove that  $n_1 = n_2$  where

$$n_1(x) = \frac{x^2 + 5x}{x^2 + 10x + 25}, n_2(x) = \frac{2x}{2x + 10}$$


---

... End of the questions ...





# Prep. 3 \_ Model (03)



## [Q1] A) Choose the correct answer:

- (1) The intersection point of two lines  $X + 2 = 0$  ,  $Y = X$  is .....  
 a)  $(2, 2)$       b)  $(2, 0)$       c)  $(-2, -2)$       d)  $(0, 0)$
- (2) If  $n(x) = \frac{x+1}{x-2}$  is an algebraic fraction, then the domain in which it has a multiplicative inverse is .....  
 a)  $R - \{2\}$       b)  $R - \{-1, 2\}$       c)  $R - \{-1\}$       d)  $\{-1, 2\}$
- (3) If the two equations  $X + 2Y = 1$  ,  $X + KY = 2$  have a one solution in  $R \times R$ , the  $K \neq$  .....  
 a) 2      b) 4      c) -2      d) -4

**B):** By using a general formula and without using calculator, find in  $R$  the solution set of the equation  $X(X - 3) = -1$ , approximating the result to one decimal place

## [Q2] A) Choose the correct answer:

- (1) If the curve of quadratic function passing through the points  $(2,0)$  ,  $(-3,0)$  ,  $(0,-6)$ , then the solution set of  $F(x) = 0$  in  $R$  is .....  
 a)  $\{-2,3\}$       b)  $\{3,2\}$       c)  $\{2,-3\}$       d)  $\{-3,-6\}$
- (2) The simplest form of  $n(x) = \frac{3-X}{X-3}$  where  $X \in R - \{3\}$  is .....  
 a) 1      b) -1      c) 3      d) -3
- (3) If A is an event in sample space of a random experiment, then  $P(A) =$  .....  
 a) 1      b) -1      c)  $1 - P(A)$       d)  $P(A) - 1$

**B):** If  $(a, 2b)$  is a solution in  $R$  of two equations  $3X - Y = 5$  ,  $X + Y = -1$ .  
 Find the value of  $a$  ,  $b$

**[Q3]**

**A)**  $n_1, n_2$  are two algebraic fractions where  $n_1(x) = \frac{x^2-4}{x^2+x-6}$ ,  $n_2(x) = \frac{x^2-x-6}{x^2-9}$

prove that  $n_1(x) = n_2(x)$  for all the values of  $x$  which belongs to the common domain and find this domain

---

**B)** Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of two equations

$$X + Y = 3, X^2 + XY = 6$$


---

**[Q4]**

**A)** Find in the simplest form:  $n(x) = \frac{x^2+3x}{x^2+2x-3} - \frac{x-2}{x^2-3x+2}$

And showing its domain

---

**B)** Find in the simplest form and showing its domain:

$$n(x) = \frac{x^3 - x^2 - 2x}{x^2 - 5x + 6} \times \frac{x^2 + 2x - 15}{x^3 + 6x^2 + 5x} \text{ then find } n(7), n(3) \text{ if possible}$$


---

**[Q5]**

**A)** If  $n_1(x) = \frac{x-a}{x+b}$ , and set of zeroes of  $n_1(x)$  is  $\{5\}$  and domain of  $n_1(x)$  is  $\mathbb{R} - \{3\}$ , find the value of  $a, b$ .

If  $n_2(x) = \frac{x-1}{x-3}$  **find**  $n_1(x) + n_2(x)$  in the simplest form

---

**B)** If  $A, B$  are two events of the sample space of a random experiment, and  $P(A) = 0.7$ ,  $P(B) = 0.6$ ,  $P(A \cap B) = 0.4$ , Find

①  $P(A \cup B)$

② The occurrence of one of the two events but not the other

---

\*\*\* End of the questions \*\*\*





# Prep. 3 - Model (04)



## [Q1] A) Choose the correct answer:

(1) The solution set of the equation  $X^2 + 4 = 0$  in  $\mathbb{R}$  is .....

- a)  $\emptyset$                       b)  $\{2\}$                       c)  $\{-2\}$                       d)  $\{2, -2\}$

(2) If  $a^2 - b^2 = 6$ ,  $a - b = \sqrt{3}$ , then  $(a + b)^2 = \dots\dots\dots$

- a)  $2\sqrt{3}$                       b)  $3\sqrt{3}$                       c)  $\sqrt{3}$                       d) 12

(3) If  $A, B$  are two mutually exclusive events, then  $P(A \cap B) = \dots\dots$

- a) Zero                      b)  $\emptyset$                       c)  $\frac{1}{6}$                       d) 1

**B):** By using a general formula and without using calculator, find in  $\mathbb{R}$  the solution set of the equation  $X^2 + 2X - 1 = 0$ , approximating the result to one decimal place.

## [Q2] A) Choose the correct answer:

(1) The set of zeroes of  $F(x) = -3x$  is .....

- a)  $\emptyset$                       b)  $\{0\}$                       c)  $\{3\}$                       d)  $\mathbb{R} - \{3\}$

(2) The simplest form of  $n(x) = \frac{3-x}{x-3}$  where  $x \neq 3$  is .....

- a) 1                      b) -1                      c) 3                      d) -3

(3) If the domain of  $n(x) = \frac{x+1}{x^2-kx+4}$  is  $\mathbb{R} - \{2\}$ , then  $K = \dots\dots\dots$

- a) 2                      b) -2                      c) 4                      d) -4

**B):** If  $n(x) = \frac{x^2 + 3x}{x^3 + 27}$

Find  $n^{-1}(x)$  in the simplest form showing the domain of  $n^{-1}(x)$

**[Q3]**

**A)** Find in the simplest form:  $n(X) = \frac{x^2 - 2x + 1}{x^3 - 1} \div \frac{x - 1}{x^2 + x + 1}$

And showing its domain

---

**B)** Find in  $\mathbf{R} \times \mathbf{R}$  the solution set of two equations

$$Y - X = 2, X^2 + XY = 4$$


---

**[Q4]**

**A)** If A , B are two events of the sample space of a random experiment,

and  $P(A) = \frac{1}{4}$  ,  $P(B) = \frac{2}{3}$  , Find  $P(A \cup B)$  if :

①  $P(A \cap B) = \frac{1}{6}$

②  $A \subset B$

---

**B)** Find in  $\mathbf{R} \times \mathbf{R}$  the solution set of two equations

$$X = Y + 4, 3X + 4Y = 5$$


---

**[Q5]**

**A)** If  $n_1(x) = \frac{x^2}{x^3 - 3x^2}$  ,  $n_2(x) = \frac{x}{x^2 - 3x}$

Prove that  $n_1 = n_2$

---

**B)** Find in the simplest form:  $n(X) = \frac{3x - 6}{x^2 - 4} - \frac{9}{2 - x - x^2}$

And showing its domain

---

••• End of the questions •••





# Prep. 3 \_ Model (05)



## [Q1] A) Choose the correct answer:

- (1) The solution set of two equations  $X - 3 = 0$  ,  $Y = 4$  in  $R \times R$  is .....  
 a)  $\{ 3 , 4 \}$       b)  $\{(3,4)\}$       c)  $\{(4,3)\}$       d)  $\emptyset$
- (2) If  $A$  ,  $B$  are two events in sample space of a random experiment,  $A \subset B$ , then  $P ( A \cup B ) = \dots\dots\dots$   
 a)  $P ( B )$       b)  $P ( A )$       c)  $P ( A \cap B )$       d) Zero
- (3) If  $3^Y \times 5^Y = 225$ , then  $Y = \dots\dots\dots$   
 a) 2      b) 15      c) Zero      d) 20

## B):

Find in  $R \times R$  the solution set of two equations:

$$3X - Y = 5 \text{ , } X + 2Y = 4$$

## [Q2] A) Choose the correct answer:

- (1) The domain of the additive inverse of  $n(x) = \frac{x+2}{x-3}$   
 a)  $R - \{3\}$       b)  $R - \{-2\}$       c)  $R - \{-2,3\}$       d)  $R$
- (2) The set of zeroes of  $F(x) = X^2 + 9$  in  $R$  is .....  
 a)  $R$       b)  $\emptyset$       c)  $\{ 3 \}$       d)  $\{ 3 , -3 \}$
- (3) The curve  $Y = a X^2 + b x + C$  cut  $Y$ -axis at the point .....  
 a)  $( 0 , b )$       b)  $( b , 0 )$       c)  $( c , 0 )$       d)  $( 0 , 7 )$

## B):

Find in the simplest form:  $n (X) = \frac{x^2+x}{x^2-1} - \frac{5-x}{x^2-x+5}$

And showing its domain

**[Q3]**

**A)** If A , B are two events of the sample space of a random experiment, and  $P(A) = 0.6$  ,  $P(B) = 0.5$  ,  $P(A \cap B) = 0.3$  , Find

①  $P(A \cup B)$

②  $P(B^c)$

---

**B)** Find in the simplest form:  $n(X) = \frac{x^3-1}{x^2-2x+1} - \frac{2x-2}{x^2+x+1}$

And showing its domain

---

**[Q4]**

**A)** Prove that  $n_1(x) = n_2$

Where  $n_1(x) = \frac{x^2-x}{x^3-2x^2}$  ,  $n_2(x) = \frac{x^2-3x+2}{x^3-4x^2+4x}$

---

**B)** By using a general formula and without using calculator, find in  $\mathbb{R}$  the solution set of the equation  $2X^2 + 4X + 1 = 0$ , approximating the result into two decimal places.

---

**[Q5]**

**A)** Find the solution set of two equations in  $\mathbb{R} \times \mathbb{R}$

$$X - Y = 0 \quad , \quad X = \frac{4}{y}$$


---

**B)** If  $n(X) = \frac{x^2-2x}{(x-2)(x^2+2)}$  Find:

①  $n^{-1}(x)$  showing its domain

② If  $n^{-1}(x) = 3$ , find the value of X.

---

••• End of the questions •••





# Prep. 3 - Model (06)



## [Q1] A) Choose the correct answer:

(1) If the two equations  $X - 3Y = 5$  ,  $2X + KY = 10$  have infinite number of solution, then  $K = \dots\dots\dots$

- a) 10                      b) 6                      c) -6                      d) 3

(2) If  $F(x) = X^3 - m$  ,  $Z(F) = \{3\}$  , then  $m = \dots\dots\dots$

- a) 9                      b) 27                      c) 3                      d)  $\sqrt[3]{3}$

(3) If  $AB = 3$  ,  $AB^2 = 9$ , then  $A^2B = \dots\dots\dots$

- a) 3                      b) 9                      c)  $\frac{1}{3}$                       d)  $\frac{1}{9}$

**B):** Find  $n(x)$  in the simplest form showing its domain

$$n(x) = \frac{x^2 - 12x + 36}{x^2 - 6x} \div \frac{36 - x^2}{4x + 24}$$

## [Q2] A) Choose the correct answer:

(1) If the probability that a student in exam is succeeded =  $\frac{4}{5}$ , then the probability his failed is  $\dots\dots\dots$

- a) 10 %                      b) 20 %                      c) Zero                      d) 1

(2) If the domain of  $F(x) = \frac{1}{x} - \frac{5}{x+k}$  is  $R - \{0, 3\}$  , then  $K = \dots\dots\dots$

- a) 3                      b) 6                      c) 5                      d) -3

(3) If  $X$  is a negative number, then the greatest one of the following is  $\dots\dots\dots$

- a)  $7X$                       b)  $7 + X$                       c)  $7 - X$                       d)  $\frac{7}{x}$

**B):** If the perimeter of rectangle is 14 cm, and its area is  $12 \text{ cm}^2$ . Find the length of its dimensions.

**[Q3]**

**A)** If  $n_1(x) = \frac{x-1}{x}$ ,  $n_2(x) = \frac{x^2-1}{x^2+x}$

Show that if  $n_1 = n_2$  or not? Give reason

**B)** If A , B are two events of the sample space of a random experiment, and  $P(A) = 0.3$ ,  $P(B) = m$ ,  $P(A \cup B) = 0.7$ , Find the value of m if:

①  $P(A \cap B) = 0.2$

② A , B are two mutually exclusive events

**[Q4]**

**A)** If  $n(x) = \frac{x^2-5x}{(x-5)(x^2+1)}$

① Find  $n^{-1}(x)$  showing its domain

② If  $n^{-1}(x) = 2$ , find the value of X

**B)** A point moves on the straight line  $5X - 2Y = 1$  where its Y-coordinate is twice the square of its X-coordinate. Find the coordinate of this point

**[Q5]**


**A)** In the opposite figure:

The curve represents  $F(x) = aX^2 + bX + c$ ,  $a \neq 0$

If the curve passes through point  $(0, 0)$

And the equation of line of symmetry  $X = 2$

And the maximum value  $Y = 2$ . Find value of a , b , c



**B)** Find  $n(x)$  in the simplest form showing its domain

$$n(x) = \frac{x^3 + x^2 - 2}{x-1} - \frac{2x-2}{1-x}$$

••• End of the questions •••





# Prep. 3 - Model (07)



## [Q1] A) Choose the correct answer:

(1) If the two equations  $X + 2Y = 1$ ,  $2X + KY = 2$  have one solution, the  $K \neq$  .....

- a) 1                      b) 2                      c) 4                      d) -4

(2) If the domain of  $n_1(x) = \frac{5}{x-8}$  equal the domain of  $n_2(x) = \frac{x-3}{x+k}$ , then  $K =$  .....

- a) 8                      b) -8                      c) 24                      d) -3

(3) Twice a number formed from two digits, its units  $Y$  and tens  $X$  is

- a)  $2Y + 10X$       b)  $2Y + 20X$       c)  $2X + 10Y$       d)  $2x + 20Y$

**B):** By using a general formula and without using calculator, find in  $R$  the solution set of the equation  $\frac{5}{x^2} - \frac{2}{x} = 1$ , and  $\sqrt{6} \simeq 2.45$

## [Q2] A) Choose the correct answer:

(1) A bag contains **20** cards numbered from **1** to **20**, on card is chosen randomly, the probability of chosen card has a number Divisible by **2** and **3** together = .....

- a)  $\frac{1}{2}$                       b)  $\frac{6}{20}$                       c)  $\frac{3}{20}$                       d)  $\frac{13}{20}$

(2) The set of zeroes of  $F(x) = \frac{x^2-x+2}{x^2-4}$  in  $R$  is .....

- a)  $\{2\}$                       b)  $\{-1\}$                       c)  $\{-1, 2\}$                       d)  $\{-2, 2\}$

(3) If  $X^2 + Y^2 = 2XY$ , then  $X - Y =$  .....

- a)  $\sqrt{2XY}$                       b)  $\sqrt{2}$                       c) Zero                      d)  $\pm 1$

**B):**

If the domain of  $F(x) = \frac{b}{x} + \frac{9}{x+a}$  is  $R - \{0, 4\}$  and  $F(5) = 2$ , find the value of  $a, b$

**[Q3]**

- A)** Prove that  $n_1(x) = n_2(x)$  for all values of  $x$  which belongs to the common domain and find this domain.

$$\text{Where } n_1(x) = \frac{x^2-4}{x^2+x-6}, \quad n_2(x) = \frac{x^3-x^2-6x}{x^3-9x}$$

- B)** If  $A, B$  are two events of the sample space of a random experiment, and  $P(A \cap B) = 0.2$ ,  $P(A - B) = 0.3$ ,  $P(B - A) = 0.4$ , Find

①  $P(B)$

②  $P(A \cup B)$

**[Q4]**

- A)** Find in the simplest form:  $n(x) = \frac{x^2+3x+9}{x^3-27} + \frac{x^2-x-12}{9-x^2}$

And showing its domain

- B)** If  $(1, 2)$  is a solution of two equations, find the value of  $a, b$

Where  $aX + bY + 5 = 0$ ,  $2aX + bY - 2 = 0$

**[Q5]**

- A)** If  $n(x) = \frac{x^2-3x}{x^2-5x+6}$  Find:

①  $n^{-1}(x)$  in simplest form and showing its domain

② If  $n^{-1}(x) = 2$ , find the value of  $x$ .

- B)** Find in the simplest form:  $n(x) = \frac{x^2+2x-3}{x^2+5x+6} \div \frac{x^2+x-2}{x^2-4}$

And showing its domain then find the value of  $x$  when  $n(x) = 3$

••• End of the questions •••





# Prep. 3 - Model (08)



## [Q1] A) Choose the correct answer:

(1) If  $X = -3$  is a solution of equation  $X^2 + mX - 9 = 0$ , then  $m = \dots$

- a) 3                      b) -3                      c) Zero                      d) -9

(2) The domain of additive inverse of  $n(x) = \frac{x}{x-3}$  is .....

- a)  $\mathbb{R}$                       b)  $\mathbb{R} - \{0\}$                       c)  $\mathbb{R} - \{3\}$                       d)  $\mathbb{R} - \{0, 3\}$

(3) Number of solution of two equations  $X - \frac{1}{2}Y = 4$ ,  $2X - Y = 2$  in  $\mathbb{R}^2$  is ..... solution

- a) One                      b) Two                      c) Infinite                      d) Zero

**B):** By using a general formula and without using calculator, find in  $\mathbb{R}$  the solution set of the equation  $X + \frac{4}{X} = 6$ , approximating the result into three decimal places.

## [Q2] A) Choose the correct answer:

(1) If  $A$  is an event in sample space of a random experiment, and  $P(A) = 4P(A^c)$ , then  $P(A) = \dots$

- a) 0.8                      b) 0.6                      c) 0.4                      d) 0.2

(2) If the set of zeroes of  $F : F(x) = aX + 6$  is  $\{-2\}$ , then  $a = \dots$

- a) 3                      b) 2                      c) -2                      d) -3

(3) If  $Y = 1 - X$ ,  $(X + Y)^2 + Y = 5$ , then  $Y = \dots$

- a) 5                      b) 4                      c) 3                      d) -4

**B):**

The area of rectangle is  $77 \text{ cm}^2$ , if its length decrease by 2 cm and the width increase by 2 cm it will be square, find the area of the square.

**[Q3]**

**A)** If the domain of  $F : F(x) = \frac{x}{x^2 - 5x + m}$  is  $\mathbb{R} - \{2, c\}$

Find the value of  $m, c$

\_\_\_\_\_

**B)** If  $A, B$  are two events of the sample space of a random experiment, and  $P(B) = \frac{1}{3}$ ,  $P(A - B) = \frac{1}{4}$ , Find  $P(A)$  if .....

①  $P(A \cap B) = \frac{1}{12}$

②  $B \subset A$

**[Q4]**

**A)** If  $n(X) = \frac{x^2 - 2x}{x^4 + 3x^3 + 2x^2} - \frac{4 - x^2}{x^2 + x - 2}$

① Find  $n(x)$  in the simplest form showing its domain

② Find the solution set of  $n(x) = 0$

\_\_\_\_\_

**B)** If  $F(x) = aX^2 + b$  and  $F(1) = 5$ ,  $F(2) = 11$

Find the value  $F(4)$

\_\_\_\_\_

**[Q5]**

**A)** Prove that  $n_1(x) = n_2(x)$  for all values of  $X$  which belongs to the common domain and find this domain.

Where  $n_1(x) = \frac{x^2 + x - 12}{x^2 + 5x + 4}$ ,  $n_2(x) = \frac{x^2 - 2x - 3}{x^2 + 2x + 1}$

\_\_\_\_\_

**B)** Find in the simplest form:  $n(X) = \frac{x^2 - 2x - 15}{x^2 - 9} \div \frac{x^2 - 25}{x^2 - 3x}$

And show its domain, and then find the value of  $a$  if  $n(a) = \frac{1}{3}$

\_\_\_\_\_

\*\*\* End of the questions \*\*\*





# Prep. 3 - Model (09)



## [Q1] A) Choose the correct answer:

(1) The two straight line  $3X + 5Y = 0$  ,  $5X - 3Y = 0$  intersecting at ....

- a) Origin point   b) First quad.   c) Second quad.   d) Fourth quad.

(2) The additive inverse of the fraction  $\frac{x+7}{x-5}$  is .....

- a)  $\frac{7-x}{x+5}$       b)  $\frac{x+7}{5-x}$       c)  $\frac{-(x+7)}{5-x}$       d)  $\frac{x-7}{5-x}$

(3) If A is an event in a sample space of a random experiment, and  $2P(A) = 3P(A^c)$ , then  $P(A) = \dots\dots\dots$

- a) 0.8      b) 0.6      c) 0.4      d) 0.2

**B):** By using a general formula and without using calculator, find in  $\mathbb{R}$  the solution set of the equation  $\frac{1}{x} + \frac{8}{x^2} = 1$  , approximating the result into three decimal places.

## [Q2] A) Choose the correct answer:

(1) In the equation  $aX^2 + bX + C = 0$ , if  $b^2 - 4ac < 0$  , then the number of roots of the equation in  $\mathbb{R}$  is .....

- a) 1      b) 2      c) Zero      d) Infinite

(2) If  $n(x) = \frac{x-1}{x+2}$  , then  $n^{-1}(4) = \dots\dots\dots$

- a) -1      b) Zero      c) 3      d) Undefined

(3) If  $X^2 - Y^2 = 6$  ,  $X - Y = \sqrt{3}$  , then  $(X + Y)^2 = \dots\dots\dots$

- a)  $2\sqrt{3}$       b)  $3\sqrt{3}$       c)  $\sqrt{3}$       d) 12

**B):**

If the length of diagonal in a rectangle is 5 cm, and its perimeter is 14 cm. find its area?

**[Q3]**

**A)** If the set of zeroes of  $F: F(x) = \frac{ax^2-6x+8}{bx-4}$  is  $\{4\}$ , and its domain is  $R - \{2\}$ , find the value of  $a, b$ .

**B)** If  $A, B$  are two events of the sample space of a random experiment, and  $P(A) = \frac{2}{5}$ ,  $P(A \cup B) = \frac{4}{5}$ ,  $P(B) + 7P(A \cap B) = 2$   
Find ①  $P(B)$  ②  $P(B - A)$

**[Q4]**


**A)** If  $n(x) = \frac{x^2-4}{x^2+x-2} + \frac{5-10x}{3x-1-2x^2}$   
Find  $n(x)$  in the simplest form showing its domain

**B) In the opposite figure:**

If the equation of Straight line  $L_1$  is  $Y = 2X$ ,

And equation of  $L_2$  is  $X + Y = 6$  where  $L_1 \cap L_2 = \{b\}$

$O$  is origin point,  $A \in \overleftrightarrow{XX'}$ . Find the area of  $\triangle OAB$


**[Q5]**

**A)** Prove that  $n_1(x) = n_2(x)$

$$\text{Where } n_1(x) = \frac{x^2-x}{x^3-2x^2}, \quad n_2(x) = \frac{x^2-3x+2}{x^3-4x^2+4x}$$

**B)** Find in the simplest form:  $n(x) = \frac{x^2+x-6}{x^2+5x+6} \div \frac{x^3-2x^2+x-2}{x^3+2x^2+x+2}$   
And show its domain.

\*\*\* End of the questions \*\*\*





# Prep. 3 - Model (10)



## [Q1] A) Choose the correct answer:

(1) If the two equations  $X + 4Y = m$  ,  $3X + KY = 21$  have infinite number of solution in  $R \times R$ , then  $K + m = \dots\dots\dots$

- a) 19                      b) 20                      c) 21                      d) 22

(2) The common domain of two fractions  $\frac{2}{x^2-1}$  ,  $\frac{5x}{x^2-x}$  is  $\dots\dots\dots$

- a)  $R - \{1\}$               b)  $R - \{0,1\}$               c)  $R - \{-1,1\}$               d)  $R - \{0,-1,1\}$

(3) If a coin is throwing once, the probability of appear a tail is  $\dots\dots\dots$

- a) 100 %              b) 50 %              c) 25 %              d) Zero

**B):** By using a general formula and without using calculator, find in  $R$  the solution set of the equation  $\frac{x^2}{9} + \frac{4}{3}X = -2$  , approximating the result into three decimal places.

## [Q2] A) Choose the correct answer:

(1) If the solution set of the equation  $4X^2 + 4X + C = 0$  in  $R$  is  $\{-\frac{1}{2}\}$  then the value of  $C = \dots\dots\dots$

- a) 2                      b) 1                      c) -1                      d) -8

(2) If  $n(x) = \frac{x^2-x}{x^2-1}$  ,  $n^{-1}(K) = 3$  , then  $K = \dots\dots\dots$

- a)  $-\frac{1}{2}$                       b)  $\frac{1}{2}$                       c)  $\frac{3}{4}$                       d)  $1\frac{1}{3}$

(3) If domain of  $F(x) = \frac{x+b}{x+a}$  is  $R - \{-2\}$  , and  $F(0) = 3$ , then  $a + b = \dots\dots\dots$

- a) 2                      b) 6                      c) 8                      d) 10

## **B):**

Find in  $R \times R$  the solution set of two equations

$$X + Y = 2 \quad , \quad \frac{1}{x} + \frac{1}{y} = 2 \quad \text{where } X \neq 0, Y \neq 0$$

**[Q3]**

**A)** If  $n_1(x) = \frac{x}{x+a}$ ,  $n_2(x) = \frac{x^3+bx}{x^3+ax^2+x+5}$  and  $n_1 = n_2$ . Find the value of  $a$ ,  $b$

**B)** If  $A$ ,  $B$  are two events of the sample space of a random experiment, and  $P(A) = 0.6$ ,  $P(B) = 0.7$ ,  $P(A \cap B) = 0.4$

Find ① The probability of non-occurrence of  $A$ ,  $B$  together.

② The probability of occurrence at least one of them.

**[Q4]**

**A)** Find in the simplest form:  $n(x) = \frac{x-6}{2x^2-15x+18} + \frac{x-5}{15-13x+2x^2}$


And show its domain,

**B) In the opposite figure:**

If the equation of Straight line  $L_1$  is  $Y = 3X$ ,

And equation of  $L_2$  is  $X + Y = 8$  where  $L_1 \cap L_2 = \{b\}$

$O$  is origin point,  $A \in \overleftrightarrow{YY'}$ . Find the area of  $\triangle OAB$


**[Q5]**


**A)** Find in the simplest form:  $n(x) = \frac{x^2-2x-15}{x^2-9} \div \frac{2x-10}{x^2-6x+9}$

And show its domain.

**B)** If  $n(x) = ax - 3$ ,  $F(x) = a^2x^2 - 12x + 9$  and  $Z(n) = Z(f)$

Find the value of  $a$  and  $Z(F)$ .

... End of the questions ...



Answer the following questions :

**1 Choose the correct answer :**

- (1) The set of zeroes of the function  $f$  : where  $f(x) = -3x$  is .....
- (a)  $\{0\}$  (b)  $\{3\}$  (c)  $\{-3\}$  (d)  $\mathbb{R} - \{3\}$
- (2) If  $A \subset S$  of a random experiment ,  $P(A) = P(\bar{A})$  , then  $P(A) = \dots\dots\dots$
- (a) 1 (b)  $\frac{1}{2}$  (c)  $\frac{1}{4}$  (d)  $\frac{1}{8}$
- (3) If  $x$  is a negative number, then the greatest number of the following is .....
- (a)  $5x$  (b)  $\frac{5}{x}$  (c)  $5 + x$  (d)  $5 - x$
- (4) The domain of the function  $f : f(x) = \frac{x-3}{4}$  is .....
- (a)  $\mathbb{R}$  (b)  $\mathbb{R} - \{-4\}$  (c)  $\mathbb{R} - \{-4, 3\}$  (d)  $\emptyset$
- (5) If the sum of ages of a father and his son now is 47 years , then the sum of their ages after 10 years = ..... years.
- (a) 27 (b) 37 (c) 57 (d) 67
- (6) If the two equations  $x + 2y = 1$  ,  $2x + ky = 2$  has only one solution , then  $k \neq \dots\dots\dots$
- (a) 1 (b) 2 (c) 4 (d) -4

**2 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two following equations algebraically :**

$$x + 3y = 7 \quad , \quad 5x - y = 3$$

**[b] Find  $n(x)$  in its simplest form , showing the domain of  $n$  :**

$$n(x) = \frac{x^2 + x}{x^2 - 1} - \frac{x + 5}{x^2 + 4x - 5}$$

**3 [a] Find in  $\mathbb{R}$  the solution set of the following equation by using the general rule :**

$$x^2 - 4x + 1 = 0 \text{ rounding the results to two decimal places.}$$

**[b] If  $n_1(x) = \frac{2x}{2x+6}$  ,  $n_2(x) = \frac{x^2+3x}{x^2+6x+9}$  , then prove that :  $n_1 = n_2$**



- 4 [a] If A and B are two events from a sample space of a random experiment , and  
 $P(A) = 0.7$  ,  $P(B) = 0.6$  ,  $P(A \cap B) = 0.4$  , then find :  
 (1)  $P(A \cup B)$  (2)  $P(A - B)$

[b] Find  $n(X)$  in its simplest form , showing the domain of  $n$  :

$$n(X) = \frac{X^3 - 8}{X^2 - 3X + 2} \times \frac{X + 1}{X^2 + 2X + 4}$$

- 5 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two following equations :

$$X - y = 1 \quad , \quad X^2 - y^2 = 25$$

[b] If  $n(X) = \frac{X^2 - 3X}{(X - 3)(X^2 + 2)}$

, then find :  $n^{-1}(X)$  in the simplest form , showing the domain of  $n^{-1}$  .

## 2 Alexandria Governorate



Answer the following questions :

- 1 Choose the correct answer from those given ones :

(1) If A , B are two mutually exclusive events ,  $P(B) = 0.5$  and  $P(A \cup B) = 0.7$   
 , then  $P(A) = \dots\dots\dots$

- (a) 0.02 (b) 0.2 (c) 0.5 (d) 0.13

(2)  $(X + 1)^2 = \dots\dots\dots$

- (a)  $X^2 + 1$  (b)  $X^2 - 1$  (c)  $X^2 - X + 1$  (d)  $X^2 + 2X + 1$

(3) The additive inverse of the fraction  $\frac{3}{X^2 + 1}$  is  $\dots\dots\dots$

- (a)  $-\frac{3}{X^2 + 1}$  (b)  $\frac{X^2 + 1}{3}$  (c)  $\frac{X^2 + 1}{-3}$  (d)  $\frac{3}{X^2 - 1}$

(4) If  $X$  is a negative real number , then the greatest number of the following numbers  
 is  $\dots\dots\dots$

- (a)  $3 + X$  (b)  $3X$  (c)  $3 - X$  (d)  $\frac{3}{X}$

(5) If  $X = 2$  and  $y = 3$  , then  $(y - 2X)^{10} = \dots\dots\dots$

- (a) 10 (b) -1 (c) -10 (d) 1

(6) The point of intersection of the two straight lines  $X = 2$  and  $X + y = 6$  is  $\dots\dots\dots$

- (a) (2 , 6) (b) (2 , 4) (c) (4 , 2) (d) (6 , 2)

- 2 [a] If A and B are two events of the sample space (S) of a random experiment such that :  
 $P(A) = 0.7$  ,  $P(A \cap B) = 0.3$  Find :  $P(A - B)$

[b] Find  $n(X)$  in the simplest form showing the domain of  $n$  , where :

$$n(X) = \frac{X^2 + 2X + 4}{X^3 - 8} - \frac{9 - X^2}{X^2 + X - 6}$$

- 3 [a] Find the common domain of  $n_1$  ,  $n_2$  to be equal such that :

$$n_1(X) = \frac{X^2 + 3X + 2}{X^2 - 4} , n_2(X) = \frac{X^2 - 1}{X^2 - 3X + 2}$$

[b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :  $X + y = 7$  ,  $X^2 + y^2 = 25$

- 4 [a] Find  $n(X)$  in the simplest form showing the domain of  $n$  , where :

$$n(X) = \frac{X}{X-2} \div \frac{X+3}{X^2 - X - 2}$$

[b] Find in  $\mathbb{R}$  the solution set of the equation :  $3X^2 - 5X - 4 = 0$

, by using the general rule , rounding the result to two decimal places.


- 5 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations graphically :

$$X + y = 4 , 2X - y = 2$$

[b] If set of zeroes of the function  $f : f(X) = aX^2 + X + b$  is  $\{0, 1\}$

find the value of each two constants a and b

### 3 El-Kalyoubia Governorate



Answer the following questions :

- 1 Choose the correct answer :

(1) Twice the number  $X$  subtracted by 3 is .....

- (a)  $X - 3$  (b)  $2X + 3$  (c)  $2X - 3$  (d)  $3 - 2X$

(2) The domain of the function  $f$  where  $f(X) = \frac{X+2}{5X}$  is .....

- (a)  $\mathbb{R} - \{5\}$  (b)  $\mathbb{R} - \{-5\}$  (c)  $\mathbb{R}$  (d)  $\mathbb{R} - \{\text{zero}\}$

(3) If  $P(A) = 4P(\bar{A})$  , then  $P(A) =$  .....

- (a) 0.8 (b) 0.6 (c) 0.4 (d) 0.2

(4) If  $X$  is a negative number , then the greatest number of the following is .....

- (a)  $5 - X$  (b)  $5 + X$  (c)  $\frac{5}{X}$  (d)  $5X$



(5) If  $2^7 \times 3^7 = 6^k$ , then  $k = \dots\dots\dots$

- (a) 14 (b) 7 (c) 6 (d) 5

(6) If  $x^2 - y^2 = 2(x + y)$  where  $(x + y) \neq \text{zero}$ , then  $(x - y) = \dots\dots\dots$

- (a) 2 (b) 4 (c) 6 (d) 8

2 [a] If  $n(x) = \frac{x^3 - 8}{x^2 - x - 2} \div \frac{x^2 + 2x + 4}{2x^2 - x - 3}$

Find  $n(x)$  in its simplest form showing the domain of  $n$

[b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :

$$2x = 1 - y, \quad x + 2y = 5 \text{ in } \mathbb{R} \times \mathbb{R}$$

3 [a] If  $A, B$  are two events in a random experiment,  $P(A) = 0.7$ ,  $P(B) = 0.6$  and  $P(A \cap B) = 0.4$

Find : (1)  $P(A \cup B)$  (2)  $P(A - B)$

[b] Find the solution set of the two equations :  $y - x = 3$ ,  $x^2 + y^2 - xy = 13$  in  $\mathbb{R}^2$

4 [a] If  $n(x) = \frac{x^2 + x}{x^2 - 1} - \frac{x - 5}{x^2 - 6x + 5}$  Find  $n(x)$  in its simplest form, showing the domain of  $n$

[b] By using the formula, find in  $\mathbb{R}$  the solution set of the equation :  $x^2 - 2x - 6 = 0$   
(Approximate to the nearest one decimal)

5 [a] If  $n_1(x) = \frac{x^2 + 2x}{x^2 + 4x + 4}$ ,  $n_2(x) = \frac{2x}{2x + 4}$ , prove that :  $n_1 = n_2$

[b] If  $n(x) = \frac{x - 2}{x + 1}$

Find : (1) The domain of  $n^{-1}$  (2)  $n^{-1}(3)$

4

El-Sharkia Governorate



Answer the following questions : (Calculator is allowed)

1 Choose the correct answer from those given :

(1) In the experiment of rolling a regular die once, the probability of appearance of an even number on the upper face = .....

- (a)  $\frac{1}{6}$  (b)  $\frac{1}{3}$  (c)  $\frac{1}{2}$  (d)  $\frac{5}{6}$



(2) The set of zeroes of the function  $f : f(x) = x^2 + 1$  is .....

- (a)  $\{1\}$  (b)  $\{-1\}$  (c)  $\{-1, 1\}$  (d)  $\emptyset$

(3) The point of intersection of the two straight lines  $x + 2 = 0$  and  $y - 3 = 0$  is .....

- (a)  $(-2, -3)$  (b)  $(-2, 3)$  (c)  $(2, -3)$  (d)  $(2, 3)$

(4) If  $2^5 \times 3^5 = m \times 6^4$ , then  $m =$  .....

- (a) 1 (b) 2 (c) 3 (d) 6

(5) The domain of the multiplicative inverse of the algebraic fraction  $\frac{x+2}{x+5}$  is .....

- (a)  $\mathbb{R}$  (b)  $\mathbb{R} - \{-5\}$  (c)  $\mathbb{R} - \{-2\}$  (d)  $\mathbb{R} - \{-2, -5\}$

(6) If  $(7^{a-2}, 3) = (1, b+5)$ , then  $a + b =$  .....

- (a) -1 (b) zero (c) 1 (d) 2

**2** [a] By using the general rule solve in  $\mathbb{R}$  the equation :  $x(x-1) = 4$  taking  $\sqrt{17} \approx 4.12$

[b] If A and B are two events in a sample space for a random experiment, and if

$$P(A) = 0.8, \quad P(B) = 0.7 \text{ and } P(A \cap B) = 0.6$$

**Find :** (1) The probability of non occurrence of the event A

(2) The probability of occurrence one of the two events at least.

**3** [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :  $x - y = 4$ ,  $3x + 2y = 7$

[b] If  $n_1(x) = \frac{x^2 - 3x + 9}{x^3 + 27}$ ,  $n_2(x) = \frac{2}{2x + 6}$  **Prove that :**  $n_1 = n_2$

**4** [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :  $x - y = 1$ ,  $x^2 - y^2 = 5$


[b] Find  $n(x)$  in the simplest form showing the domain :

$$n(x) = \frac{x^2 + 2x + 4}{x^3 - 8} - \frac{9 - x^2}{x^2 + x - 6} \text{ and find : } n(58)$$

**5** [a] If  $n(x) = \frac{x^3 - x}{x^2 - 2x + 1} \times \frac{2x - 2}{x^2 + x}$

**Find :**  $n(x)$  in the simplest form showing the domain.

[b] If the set of zeroes of the function  $f$  where  $f(x) = \frac{ax^2 - 6x + 8}{bx - 4}$  is  $\{4\}$  and its domain is  $\mathbb{R} - \{2\}$ , then find :  $a, b$



## 5 El-Monofia Governorate

Answer the following questions :

### 1 Choose the correct answer :

(1) If  $a < \sqrt{3} < b$ , then  $(a, b)$  is .....

- (a)  $(0, 1)$  (b)  $(2.5, 3.5)$  (c)  $(1, 2)$  (d)  $(2, 3)$

(2) If the curve of the quadratic function does not intersect the  $X$ -axis at any point, then the number of solutions of the equation  $f(X) = 0$  in  $\mathbb{R}$  is .....

- (a) zero (b) one solution. (c) two solutions. (d) an infinite number.

(3) If  $2^8 \times 3^8 = X \times 6^8$ , then  $X =$  .....

- (a) 2 (b) 3 (c) 6 (d) 1

(4) The set of zeroes of the function  $f : f(X) = \frac{X^2 - 9}{X - 3}$  is .....

- (a)  $\{3\}$  (b)  $\{-3\}$  (c)  $\{3, -3\}$  (d)  $\emptyset$

(5) If  $f(X) = 6X^2 + 3X(1 - 2X)$  is a polynomial function, then its degree is .....

- (a) first. (b) second. (c) third. (d) fourth.

(6) If  $A$  and  $B$  are two mutually exclusive events of random experiment then :

$P(A \cap B) =$  .....

- (a)  $P(A \cup B)$  (b)  $P(A) + P(B)$  (c)  $\emptyset$  (d) zero

### 2 [a] If $(2a + b, 3) = (18, a - b)$ :

Find the value of  $a$  and  $b$  (Indicating the steps of the solution).

[b] By using the general formula, find in  $\mathbb{R}$  the solution set for the following equation :

$$(X - 4)(X - 2) = 1 \text{ (knowing that : } \sqrt{2} \approx 1.41)$$

### 3 [a] If the domain of the function $n$ where : $n(X) = \frac{4}{X + a} + \frac{b}{2X}$

is  $\mathbb{R} - \{0, -5\}$  and  $n(3) = 1$ , find the values of  $a$  and  $b$

[b] Find  $n(X)$  in the simplest form showing the domain where :

$$n(X) = \frac{X^2 + 4X + 3}{X - 1} \div \frac{X^2 + 3X}{X^2 - X}$$

### 4 [a] Find $n(X)$ in the simplest form showing the domain where :

$$n(X) = \frac{X^2 + X + 1}{X^4 - X} + \frac{X + 3}{3 - 2X - X^2} \text{ and if } n(a) = -2, \text{ find the value of } a$$



[b] A right angled triangle in which the length of one of the sides of right angled is 5 cm. and its perimeter is 30 cm. Find the area of the triangle.

(Indicating the steps of the solution).

5 [a] If  $n_1(x) = \frac{x^2 - 4}{x^2 + x - 6}$  ,  $n_2(x) = \frac{x^3 - x^2 - 6x}{x^3 - 9x}$

**Prove that :**  $n_1(x) = n_2(x)$  for all values of  $x$  which belong to the common domain and find this domain.

[b] If A and B are two events of the sample space of a random experiment


$$P(A) = \frac{5}{9} , P(B) = \frac{2}{9} , P(A \cap B) = \frac{1}{9}$$

**Find :** (1)  $P(A \cup B)$

(2) The probability of non occurrence any of the two events.

(3) The probability of occurrence of event A only.

## 6 El-Gharbia Governorate



*Answer the following questions :*

1 Choose the correct answer from those given :

(1) If the solution set of the equation  $x^2 - ax + 4 = 0$  is  $\{-2\}$  , then  $a = \dots\dots\dots$

- (a) -2 (b) -4 (c) 2 (d) 4

(2) If  $n(x) = \frac{x+2}{x-5}$  , then the domain of  $n^{-1}$  is  $\dots\dots\dots$

- (a)  $\{2, -5\}$  (b)  $\{-2, 5\}$  (c)  $\mathbb{R} - \{-2, 5\}$  (d)  $\mathbb{R} - \{2, -5\}$

(3) If A and B are two mutually exclusive events of a random experiment

, if  $P(A) = \frac{1}{3}$  ,  $P(A \cup B) = \frac{7}{12}$  , then  $P(B) = \dots\dots\dots$

- (a)  $\frac{1}{4}$  (b)  $\frac{1}{3}$  (c)  $\frac{1}{2}$  (d) 1

(4) The set of zeroes of the function  $f : f(x) = \frac{x^2 - x - 2}{x^2 + 4}$  is  $\dots\dots\dots$

- (a)  $\{2, -2\}$  (b)  $\{-2, -1\}$  (c)  $\{2, -1\}$  (d)  $\{1, -1\}$

(5) The point of intersection of the two straight lines :  $y = 2$  ,  $x + y = 6$  is  $\dots\dots\dots$

- (a) (4 , 2) (b) (2 , 4) (c) (2 , 2) (d) (4 , 4)

(6) If the curve of the function  $f : f(x) = x^2 - x + c$  passing through the point (2 , 1) , then  $c = \dots\dots\dots$

- (a) 2 (b) 1 (c) -2 (d) -1



- 2 [a] Find in  $\mathbb{R}$  the solution set of the following equation , using the general rule , rounding the results to two decimal places :  $X(X-1) = 4$

[b] Find :  $n(X) = \frac{X^3 - 8}{X^2 + X - 6} \times \frac{X + 3}{X^2 + 2X + 4}$  in the simplest form showing the domain.

- 3 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :  $y - X = 2$  and  $X^2 + Xy - 4 = 0$

[b] Find  $n(X)$  in the simplest form , showing the domain where :  $n(X) = \frac{3}{X+1} + \frac{2X+1}{1-X^2}$

- 4 [a] Draw the graphical representation of the function  $f(X) = X^2 - 2X - 3$  in the interval  $[-2, 4]$  and from the drawing , find the solution set of the equation  $X^2 - 2X - 3 = 0$

[b] Find  $n(X)$  in the simplest form , showing the domain of  $n$  where :

$$n(X) = \frac{X^2 - 2X + 1}{X^3 - 1} \div \frac{X - 1}{X^2 + X + 1}$$

- 5 [a] If  $n(X) = \frac{X^2 - 2X}{(X-2)(X^2 + 2)}$

(1) Find  $n^{-1}(X)$  in the simplest form and determine the domain of  $n^{-1}$

(2) If  $n^{-1}(X) = 3$  what is the value of  $X$ ?

- [b] If A and B are two events in the sample space of a random experiment and if

$$P(A) = 0.7, \quad P(B) = 0.6 \text{ and } P(A \cap B) = 0.4$$

Find : (1)  $P(A \cup B)$

(2) Probability occurrence of one event without the other.

7

El-Dakahlia Governorate



Answer the following questions : (Calculators are permitted)

- 1 [a] Choose the correct answer from the given answers :

(1) The point of intersection of the two straight lines :  $X + 2 = 0$  and  $y = X$  is .....

- (a) (2 , 2)      (b) (2 , 0)      (c) (-2 , -2)      (d) (0 , 0)

(2) If  $n(X) = \frac{X+1}{X-2}$  is an algebraic fraction , then the domain in which the fraction has multiplicative inverse is .....

- (a)  $\mathbb{R} - \{2\}$       (b)  $\mathbb{R} - \{-1, 2\}$       (c)  $\mathbb{R} - \{-1\}$       (d)  $\{-1, 2\}$

(3) If there is only one solution for the equation :

$x + 2y = 1$  and  $2x + ky = 2$  in  $\mathbb{R} \times \mathbb{R}$  , then k cannot equal .....

- (a) 2 (b) 4 (c) -2 (d) -4

[b] Find in  $\mathbb{R}$  the solution set of the equation  $x(x-3) = -1$  , using the general formula (approximating the results to the nearest tenth)

2 [a] Choose the correct answer from the given answers :

(1) If the curve of the quadratic function  $f$  passes through the points  $(2, 0)$  ,  $(-3, 0)$  and  $(0, -6)$  , then the solution set of the equation  $f(x) = 0$  in  $\mathbb{R}$  is .....

- (a)  $\{-2, 3\}$  (b)  $\{3, 2\}$  (c)  $\{2, -3\}$  (d)  $\{-3, -6\}$

(2) The simplest form of the function  $n : n(x) = \frac{3-x}{x-3}$  such that  $x \in \mathbb{R} - \{3\}$  is .....

- (a) 1 (b) -1 (c) 3 (d) -3

(3) If A is an event of random experiment , then  $P(A) =$  .....

- (a) 1 (b) -1 (c)  $1 - P(A)$  (d)  $P(A) - 1$

[b] If  $(a, 2b)$  is a solution for the equations  $3x - y = 5$  and  $x + y = -1$  , find the value of a and b

3 [a]  $n_1, n_2$  are two algebraic fractions such that :  $n_1(x) = \frac{x^2 - 4}{x^2 + x - 6}$  and  $n_2(x) = \frac{x^2 - x - 6}{x^2 - 9}$

Prove that :  $n_1(x) = n_2(x)$  for all values of  $x$  which belong to the common domain and find this domain.

[b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of pair of equations :  $x + y = 3$  and  $x^2 + xy = 6$

4 [a] If  $n(x) = \frac{x^2 + 3x}{x^2 + 2x - 3} \cdot \frac{x-2}{x^2 - 3x + 2}$

Find  $n(x)$  in simplest form showing the domain of  $n$  .

[b] Find  $n(x)$  in simplest form showing the domain of  $n$  , such that :

$$n(x) = \frac{x^3 - x^2 - 2x}{x^2 - 5x + 6} \times \frac{x^2 + 2x - 15}{x^3 + 6x^2 + 5x} , \text{ then find } n(7) , n(3) \text{ if possible.}$$

5 [a] If  $f_1(x) = \frac{x-a}{x+b}$  , and the set of zeroes of  $f_1$  is  $\{5\}$  , and the domain of  $f_1$  is  $\mathbb{R} - \{3\}$  , then find the values of a and b

If  $f_2(x) = \frac{x-1}{x-3}$  , then find  $f_1(x) + f_2(x)$  in the simplest form.



[b] If A and B are two events in a sample space of a random experiment and

$P(A) = 0.7$  ,  $P(B) = 0.6$  and  $P(A \cap B) = 0.4$  , then find :

(1)  $P(A \cup B)$

(2) The probability of occurrence of one of the two events but not the other.

8

Ismailia Governorate



Answer the following questions : (Calculators are permitted)

1 Choose the correct answer from those given answers :

(1) If the age of a man now is  $X$  year , then his age after 5 years from now is ..... years.

- (a)  $X - 5$  (b)  $5 - X$  (c)  $5X$  (d)  $X + 5$

(2) The set of zero is of  $f$  where  $f(X) = X(X^2 - 2X + 1)$  is .....

- (a)  $\{0, 1\}$  (b)  $\{0, -1\}$  (c)  $\{-1, 1\}$  (d)  $\{0, 1, -1\}$

(3) If  $(5, X - 4) = (y, 3)$  , then  $X + y =$  .....

- (a) 25 (b) 12 (c) 8 (d) 6

(4) Number of solutions of the two equations :  $X + y = 2$  ,  $y - 3 = 0$  together is .....

- (a) 3 (b) 2 (c) 1 (d) zero

(5) If A and B are two mutually exclusive events , then  $P(A - B) =$  .....

- (a) zero (b)  $P(A)$  (c)  $P(B)$  (d)  $P(A \cup B)$

(6) If the curve of the function  $f$  where  $f(X) = X^2 - a$  passes through the point  $(1, 0)$  , then  $a =$  .....

- (a) -2 (b) -1 (c) zero (d) 1

2 [a] Find the solution set of the following equation in  $\mathbb{R}$  :

$$X(X - 2) = 4 \quad (\text{knowing that : } \sqrt{5} \approx 2.2)$$

[b] If  $n(X) = \frac{X^2 - 2X}{X^2 - 5X + 6}$

Find :  $n^{-1}(X)$  in the simplest form showing the domain of  $n^{-1}$

3 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the following two equations (algebraically) :

$$X + y = 5 \quad , \quad X^2 + Xy = 15$$

[b] Find  $n(X)$  in the simplest form where :  $n(X) = \frac{X}{X-4} - \frac{4X+16}{X^2-16}$



- 4 [a] A classroom consists of 40 students , 30 of them succeeded in math. 24 in science and 20 in both math. and science. If a student is chosen randomly.

Find the probability that this student is :

- (1) fail in math. (2) succeeded in math. or science

- [b] Find  $n(x)$  in the simplest form showing the domain of  $n$  :

$$n(x) = \frac{x^2 - x - 2}{x^2 - 1} \div \frac{x - 5}{x^2 - 6x + 5}$$

- 5 [a] Find  $n(x)$  in the simplest form where :  $n(x) = \frac{x^2 + 2x + 4}{x^3 - 8} + \frac{1}{x + 2}$

- [b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the following two equations (graphically) :

$$y = 3x - 1, \quad x - y + 1 = \text{zero}$$

9

Suez Governorate



Answer the following questions : (Calculators are permitted)

- 1 Choose the correct answer from those given :

- (1) The set of zeroes of  $f$  where  $f(x) = (x - 1)^2(x + 2)$  is .....

- (a)  $\{1, -2\}$  (b)  $\{-1, 2\}$  (c)  $\{-1, -2\}$  (d)  $\{1, 2\}$

- (2) If  $x - y = 2$  ,  $x^2 - y^2 = 10$  , then  $x + y =$  .....

- (a) -5 (b) 2 (c) -2 (d) 5

- (3) If  $A \subset S$  of a random experiment ,  $P(A) = P(\bar{A})$  , then  $P(A) =$  .....

- (a) zero (b) 1 (c)  $\frac{1}{2}$  (d)  $\frac{1}{4}$

- (4) If  $x$  is a negative number , then the greatest number is .....

- (a)  $3 + x$  (b)  $3 - x$  (c)  $3x$  (d)  $\frac{3}{x}$

- (5) If  $x = 3$  belongs to the solution set of the equation :  $x^2 - ax - 6 = 0$  , then  $a =$  .....

- (a) 3 (b) 2 (c) 1 (d) -1

- (6) The function  $f$  where  $f(x) = \frac{x-3}{x-4}$  has additive inverse in the domain .....

- (a)  $\mathbb{R} - \{3\}$  (b)  $\mathbb{R} - \{4\}$  (c)  $\mathbb{R} - \{-4\}$  (d)  $\mathbb{R} - \{-3\}$

- 2 [a] Find the solution set in  $\mathbb{R} \times \mathbb{R}$  :  $2x - y = 7$  ,  $3x + y = 8$

(Explain your answer showing the steps solution)

- [b] Find  $n(x)$  in the simplest form showing the domain of  $n$  where :

$$n(x) = \frac{x}{x+1} + \frac{x^2}{x^3 + x^2} \text{ , then calculate } n(3)$$

- 3 [a] Find in  $\mathbb{R} \times \mathbb{R}$  algebraically the solution set of the two equations :

$$x - 1 = 0 \text{ , } x^2 + y^2 = 10$$

- [b] If the fraction  $\frac{x+2}{x^2-4}$  is the multiplicative inverse of  $\frac{x}{h}$  where  $x \notin \{2, -2\}$  ,  
then calculate  $h$

- 4 [a] Find in  $\mathbb{R}$  the solution set for the following equations by using the formula in :

$$x^2 - 3x + 1 = 0 \text{ , knowing that } \sqrt{5} = 2.24$$

- [b] If  $n_1(x) = \frac{3x}{3x+3}$  ,  $n_2(x) = \frac{x^2+x}{x^2+2x+1}$  Prove that :  $n_1 = n_2$

- 5 [a] Find  $n(x)$  in the simplest form showing the domain of  $n$  where :

$$n(x) = \frac{x^2 + 2x + 1}{2x - 8} - \frac{x - 4}{x + 1}$$

- [b] If  $A$  and  $B$  are two events from the sample of a random experiment and

$$P(A) = 0.6 \text{ , } P(B) = 0.3 \text{ , } P(A \cap B) = 0.5$$

Find : (1)  $P(A \cup B)$  (2)  $P(\bar{B})$

## 10 Port Said Governorate



Answer the following questions :

- 1 Choose the correct answer from those given :

- (1) If the two equations :  $x + 3y = 4$  ,  $x + ay = 7$  represent two parallel straight lines ,  
then  $a = \dots\dots\dots$

- (a)  $-\frac{1}{3}$  (b)  $-3$  (c)  $3$  (d)  $1$

- (2) The domain of the multiplicative inverse of the fraction :  $\frac{x-2}{x^3+27}$  is  $\dots\dots\dots$

- (a)  $\mathbb{R} - \{2\}$  (b)  $\mathbb{R} - \{-3, 2\}$  (c)  $\mathbb{R} - \{2, -3, 3\}$  (d)  $\mathbb{R} - \{3, -3\}$



(3) If  $x^2 - y^2 = 2(x + y)$  such that :  $x + y \neq 0$  ; then  $x - y = \dots\dots\dots$

- (a) 2 (b) 4 (c) 6 (d) 8

(4) If a die is tossed once , then the probability of appearance of an odd number equals .....

- (a)  $\frac{1}{3}$  (b)  $\frac{1}{2}$  (c) 1 (d) 3

(5) The degree of the equation :  $3x + 4y + xy = 5$  is .....

- (a) zero. (b) first. (c) second. (d) third.

(6) If  $2x = 1$  , then  $\frac{1}{5}x = \dots\dots\dots$

- (a)  $\frac{2}{5}$  (b)  $\frac{1}{5}$  (c)  $\frac{1}{2}$  (d)  $\frac{1}{10}$

[2] [a] Solve in  $\mathbb{R}$  the equation :  $2x(x - 5) = 1$  approximate to the nearest one decimal.

[b] Find the common domain of  $n_1(x)$  ,  $n_2(x)$  to be equal such that :

$$n_1(x) = \frac{x^2 + 9x + 20}{x^2 - 16} , \quad n_2(x) = \frac{x^2 + 5x}{x^2 - 4x}$$

[3] [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :

$$x - 2y = 0 , \quad x^2 - y^2 = 3$$


[b] If  $n(x) = \frac{x+3}{x^2+5x-14} \div \frac{x^2+3x}{2x+14}$

Find :  $n(x)$  in its simplest form , showing the domain of  $n$

[4] [a] Find  $n$  in its simplest form , showing its domain where :  $n(x) = \frac{x^2 + x}{x^2 - 1} + \frac{x - 5}{x^2 - 6x + 5}$

[b] Use the opposite Venn diagram to calculate the probability of :

- (1) Non occurrence of the event A  
(2) The occurrence of the event B only.  
(3) Occurrence of A or B




[5] [a] If  $n(x) = \frac{x^2 - 2x}{(x - 2)(x + 2)}$

- (1) Find :  $n^{-1}(x)$  (2) If  $n^{-1}(x) = 3$  what is the value of  $x$ ?

[b] Two number , if three times a number is added to twice a second number the sum is 13 and if the first number is added to three times the second number the sum is 16 , find the two number.





Answer the following questions : (Calculators are permitted)

1 Choose the correct answer from the given ones :

(1) The solution set of the equation :  $aX^2 + bX + c = 0$  ,  $a \neq 0$  graphically is the set of  $X$  coordinates of the points of intersection of the curve of the function  $f : f(X) = aX^2 + bX + c$  with the .....

- (a) y-axis (b) X-axis (c) symmetric line (d) straight line  $y = 2$

(2) If  $a b = 12$  ,  $b c = 20$  ,  $a c = 15$  ,  $a \in \mathbb{R}^+$  ,  $b \in \mathbb{R}^+$  ,  $c \in \mathbb{R}^+$  , then  $a b c = \dots\dots\dots$

- (a) 360 (b) 3600 (c) 60 (d) 36

(3) If the algebraic fraction  $\frac{X-a}{X+5}$  have a multiplicative inverse which is  $\frac{X+5}{X+3}$  , then  $a = \dots\dots\dots$

- (a) 3 (b) -5 (c) -3 (d) 5

(4)  $\sqrt{(-2)^4 + 3^2} = \dots\dots\dots + 3$

- (a)  $2^2$  (b) 2 (c) -2 (d)  $(-2)^2$

(5) If  $P(A) = P(\bar{A})$  , then  $P(A) = \dots\dots\dots$

- (a)  $\frac{1}{2}$  (b) 1 (c)  $\frac{3}{4}$  (d) 0

(6)  $X^3 - 1 = \dots\dots\dots$

- (a)  $(X^2 - 1)(X + 1)$  (b)  $(X - 1)(X^2 + 2X + 1)$   
(c)  $(X - 1)(X^2 + X + 1)$  (d)  $(X - 1)(X^2 - 2X - 1)$

2 [a] Find :  $n(X) = \frac{X-3}{X^2-7X+12} - \frac{4}{X^2-4X}$  in the simplest form showing the domain of  $n$

[b] Find the value of  $a$  and  $b$  , knowing that :  $\{(3, -1)\}$  is the solution set of the two equations :  $aX + bY - 5 = 0$  ,  $3aX + bY = 17$

3 [a] Find in  $\mathbb{R}$  the solution set for the equation  $X(X-1) = 4$  using the general rule to the nearest hundredth.

[b] Find the common domain of  $f_1$  ,  $f_2$  to be equal such that :

$$f_1(X) = \frac{X^2 + X - 12}{X^2 + 5X + 4} , f_2(X) = \frac{X^2 - 2X - 3}{X^2 + 2X + 1}$$

- 4 [a] Two acute angles in a right-angled triangle the difference between their measures is  $50^\circ$ . Find the measure of each angle.

[b] Find  $n(X)$  in the simplest form showing the domain :


$$n(X) = \frac{x^2 - 3x + 2}{x^2 - 1} \div \frac{3x - 15}{x^2 - 4x - 5}$$

- 5 [a] If A and B are two events from a sample space of a random experiment and  $P(A) = 0.4$  ,  $P(B) = 0.5$  ,  $P(A \cup B) = 0.7$

Find : (1)  $P(A \cap B)$  (2)  $P(B - A)$

[b] If  $n(X) = \frac{x + \frac{1}{x}}{4x + \frac{4}{x}}$  Find  $n(X)$  in the simplest form showing the domain.

## 12 Kafr El-Sheikh Governorate



Answer the following questions : (Calculator is allowed)

- 1 [a] Choose the correct answer from those given :

(1) If  $x = y + 1$  ,  $(x - y)^2 + y = 3$  , then  $y = \dots\dots\dots$

(a) zero (b) 1 (c) 2 (d) 3

(2) If  $a \cdot b = 3$  ,  $a \cdot b^2 = 12$  , then  $b = \dots\dots\dots$

(a) 4 (b) 2 (c) -2 (d)  $\pm 2$

(3) If  $n(X) = \frac{x-1}{x-2}$  , then the domain of  $n^{-1} = \dots\dots\dots$

(a)  $\mathbb{R}$  (b)  $\mathbb{R} - \{1\}$  (c)  $\mathbb{R} - \{2\}$  (d)  $\mathbb{R} - \{1, 2\}$

[b] Solve in  $\mathbb{R} \times \mathbb{R}$  the two simultaneous equations :

$$x - y = 1 \quad , \quad x^2 + y^2 = 25$$

- 2 [a] Choose the correct answer from those given :

(1) The probability of the impossible event equals  $\dots\dots\dots$

(a)  $\emptyset$  (b) zero (c) 1 (d) -1

(2) If the solution set of the equation :  $x^2 + m \cdot x + 9 = 0$  is  $\{-3\}$  , then  $m = \dots\dots\dots$

(a) 5 (b) 6 (c)  $\pm 6$  (d) zero

(3) If the two equations :  $x + 3y = 6$  ,  $2x + ky = 12$  have an infinite number of solution in  $\mathbb{R} \times \mathbb{R}$  , then  $k = \dots\dots\dots$

(a) 2 (b) 6 (c) 3 (d) 1



[b] Two acute angles in a right-angled triangle the difference between their measures is  $50^\circ$   
Find the measure of each angle.

3 [a] Solve in  $\mathbb{R}$  using the (general rule) the equation :  $3x^2 = 5x + 4$  approximating the result to the nearest two decimals.

[b] Find  $n(x)$  in the simplest form showing the domain of  $n$  where :

$$n(x) = \frac{3}{x+1} + \frac{2x+1}{1-x^2}$$

4 [a] If  $A, B$  are two events from a sample space of random experiment, and

$$P(B) = \frac{1}{12}, \quad P(A \cup B) = \frac{1}{3}, \quad \text{then find } P(A) \text{ if :}$$

(1)  $A$  and  $B$  are two mutually exclusive events.

(2)  $B \subset A$

[b] If  $n_1(x) = \frac{x^2}{x^3 - x^2}$ ,  $n_2(x) = \frac{x^3 + x^2 + x}{x^4 - x}$  Prove that :  $n_1 = n_2$

5 [a] If  $n(x) = \frac{x^2 - 5x}{(x-5)(x^2+1)}$


(1) Find  $n^{-1}(x)$  and identify the domain of  $n^{-1}$

(2) If  $n^{-1}(x) = 2$ , find the value of  $x$

[b] If  $n(x) = \frac{x^2 - 3x}{x^2 - 9} \div \frac{2x}{x+3}$

Find  $n(x)$  in the simplest form showing the domain of  $n$

## 13 El-Beheira Governorate



Answer the following questions : (Calculator is allowed)

1 Choose the correct answer from the given ones :

(1) If  $f(x) = 2x$ , then  $f(1) - f(-1) = \dots\dots\dots$

(a) zero

(b) 4

(c) 2

(d) -2

(2) The two straight lines :  $x + 5y = 1$ ,  $x + 5y - 8 = 0$  are  $\dots\dots\dots$

(a) parallel.

(b) coincide.

(c) intersect and non perpendicular.

(d) perpendicular.

(3) If  $n(x^2) = 9$ , then  $n(x) = \dots\dots\dots$

(a) 81

(b) 3

(c)  $\pm 3$

(d) -3



(4) If  $n(x) = \frac{x-2}{x^2-x-6}$ , then the domain of  $n^{-1}$  is .....

- (a)  $\mathbb{R} - \{2\}$  (b)  $\mathbb{R} - \{-2, 3\}$  (c)  $\mathbb{R} - \{-2, 2\}$  (d)  $\mathbb{R} - \{-2, 2, 3\}$

(5) The degree of the equation :  $3x + 4y + xy = 5$  is .....

- (a) zero. (b) first. (c) second. (d) third.

(6) A card is drawn randomly from 20 identical cards numbered from 1 to 20, then the probability that the number of the drawn card multiple of 7 is .....

- (a) 10 % (b) 15 % (c) 20 % (d) 25 %

**2 [a] Solve in  $\mathbb{R}$  the equation :  $3x^2 = 5x + 4$  approximating the result to the nearest two decimals.**

**[b] Simplify the function  $n(x)$  where :**

$$n(x) = \frac{3x}{x^2-2x} - \frac{12}{x^2-4} \text{ showing the domain of } n$$

**3 [a] If  $f(x) = \frac{x^2-9}{x+b}$ ,  $f(4) = 1$  Find : b**

**[b] If A and B are two events in a random experiment**

$$, P(A) = 0.7, P(B) = 0.6 \text{ and } P(A \cap B) = 0.4$$

**Find the probability of :**

- (1) Non occurrence of the event A  
(2) Occurrence of one of the events but not the other.

**4 [a] The sum of two rational numbers is 12, and three times the smallest number exceeds than twice the greatest number by one Find the two numbers.**

**[b] If  $n_1(x) = \frac{x^2}{x^3-x^2}$ ,  $n_2(x) = \frac{x^3+x^2+x}{x^4-x}$ , then prove that :  $n_1 = n_2$**

**5 [a] Solve in  $\mathbb{R} \times \mathbb{R}$  the two equations :  $x - y = 1$ ,  $x^2 + y^2 = 25$**

**[b] If  $f(x) = \frac{x^2-49}{x^3-8} \div \frac{x+7}{x-2}$**

**Find :  $f(x)$  in its simplest form showing the domain of  $f$**



## 14 El-Fayoum Governorate

Answer the following questions : (Calculators are permitted)

1 Choose the correct answer from the given ones :

(1)  $(2\sqrt{2})^4 = \dots\dots\dots$

- (a) 8 (b) 16 (c) 32 (d) 64

(2) If A and B are mutually exclusive events from the sample space of a random experiment , then  $P(A \cap B) = \dots\dots\dots$

- (a) 1 (b) zero (c)  $\frac{1}{2}$  (d) -1

(3) If  $X = 1$  is the solution of the equation :  $X^2 + mX + 4 = 0$  , then  $m = \dots\dots\dots$

- (a) 1 (b) -1 (c) zero (d) -5

(4) If  $2X^2 = 5$  , then  $6X^2 = \dots\dots\dots$

- (a) 5 (b) 10 (c) 15 (d) 20

(5) If  $n(X) = \frac{X}{X-1}$  , then the domain of  $n^{-1} = \dots\dots\dots$

- (a)  $\mathbb{R} - \{0\}$  (b)  $\mathbb{R} - \{1\}$  (c)  $\mathbb{R} - \{0, 1\}$  (d)  $\mathbb{R} - \{-1\}$

(6) The sum of two consecutive integers is 17 , then the smaller number of them is  $\dots\dots\dots$

- (a) 8 (b) 9 (c) 17 (d) 72

2 [a] If  $n(X) = \frac{X^2 + X}{X^2 - X - 2} - \frac{2X + 4}{X^2 - 4}$  , find  $n(X)$  in the simplest form showing the domain of  $n$

[b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :

$$y = X + 1 \quad , \quad X^2 + y^2 = 13$$

3 [a] By using the general rule find in  $\mathbb{R}$  the solution set of the equation :

$$X^2 - 5X + 3 = 0 \quad , \quad \text{approximating the result to the nearest one decimal digit.}$$

[b] Find  $n(X)$  in the simplest form showing the domain of  $n$  where :

$$n(X) = \frac{X^3 - 1}{X^2 - 2X + 1} \div \frac{X^2 + X + 1}{2X - 2}$$

4 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations graphically :

$$y = X + 1 \quad , \quad 2X + y = 7$$

[b] Find the set of zeroes of the function  $f : f(X) = \frac{X-1}{X+1}$  , then find  $f^{-1}(2)$




- 5 [a] Find the common domain of  $n_1$  and  $n_2$  to be equal such that :

$$n_1(x) = \frac{x^2 + 2x}{x^2 + 3x + 2}, \quad n_2(x) = \frac{x^2 - x}{x^2 - 1}$$

- [b] A bag contains 10 identical cards numbered from 1 to 10, one card of them is drawn randomly, calculate the probability that the number on the drawn card is :

- (1) A prime number. (2) A number divisible by 5

## 15 Beni Suez Governorate



Answer the following questions : (Calculator is allowed)

- 1 Choose the correct answer from those given :

- (1) The probability of the impossible event equals .....

- (a)  $\emptyset$  (b) 1 (c) zero (d) -1

- (2) If  $2^x = 8$ , then  $x =$  .....

- (a) zero (b) 1 (c) 2 (d) 3

- (3) If the two straight lines which represent the two equations :

$$x + 2y = 4, \quad 2x + ky = 11 \text{ are parallel, then } k = \dots\dots\dots$$

- (a) 4 (b) 1 (c) -1 (d) -4

- (4) If  $a$  is a negative number, then the greatest number is .....

- (a)  $3 + a$  (b)  $3 - a$  (c)  $3a$  (d)  $\frac{3}{a}$

- (5) The solution set of the equation :  $x^2 + 1 = 0$  in  $\mathbb{R}$  is .....

- (a)  $\{1\}$  (b)  $\{1, -1\}$  (c)  $\{-1\}$  (d)  $\emptyset$

- (6) If  $n(x) = \frac{x-1}{x+2}$ , then  $n^{-1}(1)$  is .....

- (a) -1 (b) zero (c) 3 (d) undefined.

- 2 [a] Find the set of zeroes of the function  $f : f(x) = x^3 - x$

- [b] Find in  $\mathbb{R}$  the solution set of the following equation by using the general formula :

$$x^2 - 5x + 3 = 0 \text{ approximating the result to the nearest one decimal digit.}$$

- 3 [a] Find algebraically in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :

$$x + y = 4, \quad 2x - y = 2$$

- [b] If  $A$  and  $B$  are two events from a sample space of a random experiment

$$P(A) = 0.6, \quad P(B) = 0.5 \text{ and } P(A \cap B) = 0.3$$

- Find : (1)  $P(A - B)$  (2)  $P(A \cup B)$



4 [a] If  $n_1(x) = \frac{x^2 - 2x + 4}{x^3 + 8}$  ,  $n_2(x) = \frac{3}{3x + 6}$

Prove that :  $n_1 = n_2$

[b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :


$$x - 2 = 0 \quad , \quad x^2 + xy + y^2 = 7$$

5 [a] Find  $n(x) = \frac{x^2 - 3x + 2}{x^2 - 1} \div \frac{3x - 15}{x^2 - 4x - 5}$

in the simplest form showing the domain of  $n$

[b] If the domain of the function  $n : n(x) = \frac{x - 1}{x^2 - ax + 9}$  is  $\mathbb{R} - \{3\}$   
 , then find the value of  $a$

## 16 El-Menia Governorate



Answer the following questions : (Calculator is allowed)

1 Choose the correct answer from those given :

(1)  $(-1)^{37} - (-1)^{36} = \dots\dots\dots$

- (a) -2 (b) zero (c) 1 (d) 2

(2) The degree of the function  $f : f(x) = 2x^3 + 3x^2 - 5$  is  $\dots\dots\dots$

- (a) fourth. (b) fifth. (c) third. (d) zero.

(3) If  $a + b = 7$  ,  $a^2 - b^2 = 21$  , then  $a - b = \dots\dots\dots$

- (a) -7 (b) 7 (c) -3 (d) 3

(4) The simplest form of the function  $f : f(x) = \frac{3 - x}{x - 3}$  where  $x \neq 3$  is  $\dots\dots\dots$

- (a) 3 (b) 1 (c) -1 (d) zero

(5) The number of solutions of the two equations :

$$x - \frac{1}{2}y = 4 \quad , \quad 2x - y = 1 \text{ in } \mathbb{R}^2 \text{ is } \dots\dots\dots$$

- (a) one solution (b) two solutions.  
 (c) an infinite number. (d) zero.

(6) If a die is tossed once , then the probability of appearance of a number greater than 4 is  $\dots\dots\dots$

- (a)  $\frac{2}{3}$  (b)  $\frac{1}{6}$  (c)  $\frac{1}{3}$  (d)  $\frac{1}{2}$

2 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of :

$$x + y = \text{zero} \quad , \quad 5y^2 - 4x^2 = 36$$

[b] Find  $n(x)$  in the simplest form and determine the domain of  $n$  :

$$n(x) = \frac{x-3}{x^2-7x+12} - \frac{4}{x^2-4x}$$

[3] [a] By using the general formula find in  $\mathbb{R}$  the S.S. of :  $x^2 - x - 4 = 0$  where  $\sqrt{17} \approx 4.12$

[b] If  $n_1(x) = \frac{2x}{2x+4}$  ,  $n_2(x) = \frac{x^2+2x}{x^2+4x+4}$  Prove that :  $n_1 = n_2$

[4] [a] Find  $n(x)$  in the simplest form showing the domain of  $n$  :  $n(x) = \frac{x^2+2x+1}{2x-8} \times \frac{x-4}{x+1}$

[b] If  $(-3, 1)$  is a solution for the two equations  $ax + by = 5$  ,  $3ax + by - 17 = 0$

Find :  $a, b$

[5] [a] If the domain of  $n$  :  $n(x) = \frac{l}{x} + \frac{9}{x+m}$  is  $\mathbb{R} - \{0, -2\}$  ,  $n(4) = 1$  Find :  $l, m$

[b] If  $S$  is the sample space of a random experiment where its outcomes are equal ,  $A$  and  $B$  are two events from  $S$  , if the number of outcomes that leads to the occurrence of the event  $A = 13$  and the number of all possible outcomes of the random experiment is 24 ,  $P(A \cup B) = \frac{5}{6}$  and  $P(B) = \frac{5}{12}$

Find :

- (1) The probability of occurrence of the event  $A$
- (2) The probability of occurrence of the events  $A$  and  $B$  together.

17

Assiut Governorate



Answer the following questions : (Calculator is allowed)

[1] Choose the correct answer :

(1) The solution set of the two equations :  $x = -1$  ,  $y - 1 = 0$  in  $\mathbb{R} \times \mathbb{R}$  is .....

- (a)  $\{(-1, 1)\}$  (b)  $\{(1, -1)\}$  (c)  $\{(-1, -1)\}$  (d)  $\{(1, 1)\}$

(2) The solution set of the equation :  $2x + 4 = 0$  in  $\mathbb{N}$  is .....

- (a)  $\{2\}$  (b)  $\{-2\}$  (c)  $\{0\}$  (d)  $\emptyset$

(3) The domain of the function  $f$  where  $f(x) = \frac{x-2}{x^2+1}$  is .....

- (a)  $\mathbb{R} - \{-1\}$  (b)  $\mathbb{R} - \{1, -1\}$  (c)  $\mathbb{R} - \{1\}$  (d)  $\mathbb{R}$

(4) If  $A \subset S$  ,  $P(A) = \frac{1}{3}$  , then  $P(\bar{A}) = \dots\dots\dots$

- (a)  $\frac{1}{3}$  (b)  $\frac{2}{3}$  (c)  $\frac{1}{2}$  (d)  $\frac{3}{2}$

(5)  $|-5| = \dots\dots\dots$

- (a)  $-5$  (b)  $-\frac{1}{5}$  (c)  $5$  (d)  $\frac{1}{2}$



(6) If A and B are two mutually exclusive events of a random experiment ,  
then  $P(A \cap B) = \dots\dots\dots$

- (a)  $\emptyset$  (b) 1 (c) zero (d)  $\frac{1}{2}$

**2** [a] Find alagabrically the solution set of the two equations :

$$2x - y = 3 \quad , \quad x + 2y = 4$$

[b] Find  $n(x)$  in the simplest form showing the domain of n where :

$$n(x) = \frac{x^2 + 2x + 4}{x^3 - 8} + \frac{x^2 + x - 2}{x^2 - 4}$$

**3** [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :

$$x - y = 1 \quad , \quad x^2 + y^2 = 25$$

[b] If  $n(x) = \frac{x^2 - 3x + 2}{x^2 - 1} \div \frac{3x - 15}{x^2 - 4x - 5}$

, find  $n(x)$  in the simplest form showing the domain of n

**4** [a] Find in  $\mathbb{R}$  the solution set of the equation :  $3x^2 - 5x - 1 = 0$

approximating the result to the nearest two decimals.

[b] If  $n(x) = \frac{x^2 + 3x}{x^3 + 27}$ , find  $n^{-1}(x)$  in its simplest form showing the domain of  $n^{-1}$

**5** [a] If  $n_1(x) = \frac{x^2}{x^3 - x^2}$  ,  $n_2(x) = \frac{x^3 + x^2 + x}{x^4 - x}$  Prove that :  $n_1 = n_2$

[b] A bag contains 15 identical balls numbered from 1 to 15 , one ball is chosen randomly , if the event A is getting an odd number and the event B is getting a number divisible by 5

Find :

- (1)  $P(A)$  (2)  $P(B)$  (3)  $P(A - B)$

## 18 Souhag Governorate



Answer the following questions : (Calculator is allowed)

**1** Choose the correct answer :

(1) The set of zeroes of the function  $f$  where  $f(x) = \frac{x-3}{x+2}$  is .....

- (a) {zero} (b) {3} (c) {-2} (d) {3, -2}

(2) If  $2^n = 3$  , then  $8^n = \dots\dots\dots$

- (a) 27 (b) 9 (c) 3 (d) 6

(3) If A and B are two mutually exclusive events of a random experiment , then  $P(A \cap B) = \dots\dots\dots$

- (a)  $\emptyset$  (b) 1 (c) 2 (d) zero

(4) If  $3^x + 3^x + 3^x = 9$  , then  $x = \dots\dots\dots$

- (a) 4 (b) 2 (c) 1 (d) 9


(5) If the two equations :  $x + 3y = 6$  ,  $2x + ky = 12$  have an infinit number of solutions , then  $k = \dots\dots\dots$

- (a) 1 (b) 6 (c) 3 (d) 2

(6) In the opposite figure :

The solution set of  $f : f(x) = 0$  is  $\dots\dots\dots$

- (a)  $\emptyset$  (b)  $\{3\}$   
(c)  $\{2, 3\}$  (d)  $\{2\}$



2 [a] Solve in  $\mathbb{R}$  the equation :  $2x^2 - 5x + 1 = 0$  approximating the result to the nearest two decimals.

[b] If  $n_1(x) = \frac{x^2}{x^3 - x^2}$  ,  $n_2(x) = \frac{x^3 + x^2 + x}{x^4 - x}$  , prove that :  $n_1 = n_2$

3 [a] Solve in  $\mathbb{R} \times \mathbb{R}$  the two equations :  $x - 2y = 1$  ,  $x^2 - xy = 0$

[b] Find  $n(x)$  in the simplest form showing the domain of  $n$  where :  $n(x) = \frac{x}{x+1} + \frac{2x^2}{x^3 - x}$

4 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :

$$2x + y = 1 \quad , \quad x + 2y = 5$$

[b] If  $n(x) = \frac{x^2 - 3x}{x^2 - 9} \div \frac{2x}{x+3}$  , find  $n(x)$  in its simplest form showing the domain of  $n$

5 [a] If  $n(x) = \frac{x-2}{x+1}$  ,

Find : (1)  $n^{-1}(x)$  showing the domain of  $n^{-1}$  (2)  $n^{-1}(3)$

[b] If A and B are two events in a random experiment

,  $P(A) = 0.7$  ,  $P(B) = 0.6$  and  $P(A \cap B) = 0.4$

Find : (1)  $P(A \cup B)$  (2)  $P(A - B)$





## 19 Qena Governorate

Answer the following questions : (Calculators are permitted)

### 1 Choose the correct answer :

(1) If there are infinite numbers of solutions of the two equations

$$X + 4y = 7 \quad , \quad 3X + ky = 21 \quad , \text{ then } k = \dots\dots\dots$$

- (a) 4 (b) 7 (c) 12 (d) 21

(2) One of the solutions for the two equations :  $X - y = 2$  ,  $X^2 + y^2 = 20$  is .....

- (a)  $(-4, 2)$  (b)  $(2, -4)$  (c)  $(3, 1)$  (d)  $(4, 2)$

(3) The set of zeroes of  $f$  where  $f(X) = X^2 - 2$  is .....

- (a)  $\{2\}$  (b)  $\{-2\}$  (c)  $\{\sqrt{2}, -\sqrt{2}\}$  (d)  $\emptyset$

(4) The simplest form of  $f(X) = \frac{4X^2 - 2X}{2X}$  ,  $X \neq 0$  is .....

- (a)  $4X^2$  (b)  $2X - 1$  (c)  $2X$  (d) 2

(5) If A and B are two mutually exclusive events, then  $P(A \cap B) = \dots\dots\dots$

- (a)  $\emptyset$  (b) zero (c) 0.56 (d) 1

(6) If  $A \subset B$  , then  $P(A \cup B) = \dots\dots\dots$

- (a) zero (b)  $P(A)$  (c)  $P(B)$  (d)  $P(A \cap B)$

### 2 [a] Find in $\mathbb{R} \times \mathbb{R}$ algebraically the solution set of the two equations :

$$2X - y = 3 \quad , \quad X + 2y = 4$$

[b] If  $n_1(X) = \frac{2X}{2X+4}$  ,  $n_2(X) = \frac{X^2+2X}{X^2+4X+4}$  Prove that :  $n_1 = n_2$

### 3 [a] Find in $\mathbb{R}$ the solution set of the following equation by using the general rule :

$$3X^2 = 5X - 1 \quad (\text{Rounding the results to two decimal places})$$

[b] Find  $n(X)$  in the simplest form showing the domain of  $n$  where :

$$n(X) = \frac{X^2 + X + 1}{X} \times \frac{X^2 - X}{X^3 - 1}$$

### 4 [a] Find in $\mathbb{R} \times \mathbb{R}$ algebraically the solution set of the two equations :

$$X + y = 7 \quad , \quad Xy = 12$$

[b] Find  $n(X)$  in the simplest form showing the domain of  $n$  where :

$$n(X) = \frac{3X - 4}{X^2 - 5X + 6} + \frac{2X + 6}{X^2 + X - 6}$$

5 [a] If  $n(x) = \frac{x^2 - 2x}{(x-2)(x^2 + 2)}$

(1) Find  $n^{-1}(x)$  and identify the domain.

(2) If  $n^{-1}(x) = 3$  what is the value of  $x$ ?

[b] If A and B are two events from the sample space of a random experiment and

$P(A) = 0.7$  ,  $P(A \cap B) = 0.3$  Find :  $P(A - B)$



20

Luxor Governorate

Answer the following questions :

1 Choose the correct answer :

(1) The set of zeroes of the function  $f : f(x) = x^2 + 3$  is .....

(a)  $\{0\}$

(b)  $\emptyset$

(c)  $\{3\}$

(d)  $\{3, -3\}$

(2)  $\sqrt{16+9} = 4 + \dots$

(a) 3

(b) 5

(c) 1

(d) 7

(3) If  $\bar{A}$  is the complement event of the event A in a sample space of a random experiment , then  $P(A) + P(\bar{A}) = \dots$

(a) 2

(b) 1

(c)  $\frac{1}{2}$

(d) 3

(4) If  $3^x = 1$  , then  $x = \dots$

(a) 0

(b)  $\frac{1}{3}$

(c) 1

(d) 3

(5) The point of intersection of the two straight lines :  $y = 2$  ,  $x + y = 6$  is .....

(a) (2 , 4)

(b) (2 , 6)

(c) (6 , 2)

(d) (4 , 2)

(6) If  $(5, x-4) = (y+2, 3)$  , then  $x + y = \dots$

(a) 6

(b) 8

(c) 10

(d) 12

2 [a] Find the solution set of the two equations in  $\mathbb{R}^2$  :  $x - 2y = 0$  ,  $x^2 - y^2 = 3$

[b] If  $n(x) = \frac{x^2 - 16}{x + 4}$

Find : (1)  $n^{-1}(x)$  showing the domain of  $n^{-1}$

(2)  $n^{-1}(4)$

(3)  $n(4)$

3 [a] If  $n_1(x) = \frac{2x}{2x+4}$  ,  $n_2(x) = \frac{x^2+2x}{x^2+4x+4}$  Prove that :  $n_1 = n_2$

[b] Using the general rule find in  $\mathbb{R}$  the S.S. of the equation :

$3x^2 = 5x - 1$  (given that  $\sqrt{13} \approx 3.61$ )



- 4 [a] If A , B are two events of the sample space of a random experiment and if  
 $P(B) = \frac{1}{12}$  ,  $P(A \cup B) = \frac{1}{3}$

Find P (A) in the following cases :

(1) A and B are two mutually exclusive events

(2)  $B \subset A$

[b] If  $n(X) = \frac{x^2 + x}{x^2 - 1} - \frac{5 - x}{x^2 - 6x + 5}$

Find n (X) in the simplest form showing the domain of n.


5 [a] If  $n(X) = \frac{x^3 - 1}{x^2 - 2x + 1} \times \frac{2x - 2}{x^2 + x + 1}$

Find n (X) in the simplest form showing the domain

[b] Find graphically in  $\mathbb{R} \times \mathbb{R}$  the S.S. of the two equations :

$y = x + 4$  ,  $x + y = 4$

## 21 Aswan Governorate



Answer the following questions : (Calculators are permitted)

1 Choose the correct answer from those given :

(1) If  $x + y = 5$  , then  $3x + 3y = \dots\dots\dots$

- (a) 5 (b) 3 (c) 8 (d) 15

(2) If  $\sqrt{64 + 36} = 8 + x$  , then  $x = \dots\dots\dots$

- (a) 9 (b) 6 (c) 2 (d) 10

(3) The solution set of the two equations :  $y - 5 = 0$  ,  $y + x = 0$  in  $\mathbb{R} \times \mathbb{R}$  is  $\dots\dots\dots$

- (a)  $\{(-5, 5)\}$  (b)  $\{(5, -5)\}$  (c)  $\{(0, 5)\}$  (d)  $\{(-5, 5)\}$

(4) The set of zeroes of the function  $f : f(x) = 4$  is  $\dots\dots\dots$

- (a)  $\{-4\}$  (b)  $\{\text{zero}\}$  (c)  $\emptyset$  (d)  $\{2\}$

(5) If the probability that a student succeeded is 95 % , then the probability that he does not succeed is  $\dots\dots\dots$

- (a) 20 % (b) 5 % (c) 10 % (d) zero

(6) The solution set of the equation :  $x^2 - 4x + 4 = 0$  in  $\mathbb{R}$  is  $\dots\dots\dots$

- (a)  $\{-2\}$  (b)  $\{2\}$  (c)  $\{4, 1\}$  (d)  $\emptyset$

2 [a] Find in  $\mathbb{R} \times \mathbb{R}$  algebraically the solution set of two equations :

$x + y = 4$  ,  $2x - y = 2$

[b] If  $n(x) = \frac{x-1}{x+3}$  find  $n^{-1}(x)$  and identify the domain of  $n^{-1}$

3 [a] If  $n(x) = \frac{x^4 - 3x}{x^2 - 9} \div \frac{2x}{x+3}$ , find  $n(x)$  in the simplest form showing the domain of  $n$

[b] Find in  $\mathbb{R} \times \mathbb{R}$  algebraically the solution set of the two equations :

$$x - 2y = 0, \quad x^2 - y^2 = 3$$

4 [a] If A and B are two events from a sample space of a random experiment and

$$P(A) = \frac{1}{2}, \quad P(B) = \frac{1}{3}$$

Find  $P(A \cup B)$  if :

$$(1) P(A \cap B) = \frac{1}{8}$$

(2) A and B are mutually exclusive events.

[b] If  $n(x) = \frac{x}{x^2 + 2x} + \frac{x-2}{x^2 - 4}$ , find  $n(x)$  in the simplest form showing the domain of  $n$


5 [a] By using the formula find in  $\mathbb{R}$  the solution set of the equation

$$3x^2 - 5x + 1 = 0 \text{ rounding the result to two decimal places.}$$

[b] Find the common domain in which the two functions  $n_1$  and  $n_2$  are equal where :

$$n_1(x) = \frac{x^2 + x - 12}{x^2 + 5x + 4}, \quad n_2(x) = \frac{x^2 - 2x - 3}{x^2 + 2x + 1}$$

## 22 South Sinai Governorate



Answer the following questions : (Calculator is permitted)

1 Choose the correct answer from those given :

(1) The number of solutions of the two equations :  $x + y = 5$  and  $y - 5 = 0$  is .....

- (a) zero (b) 1 (c) 2 (d) 3

(2) The point  $(-3, 4)$  lies in ..... quadrant.

- (a) fourth (b) third (c) second (d) first

(3) The range of the set of the values : 7, 3, 6, 9 and 5 equals .....

- (a) 3 (b) 4 (c) 5 (d) 6

(4)  $(-3x) \times (-5y) = \dots\dots\dots$

- (a)  $15xy$  (b)  $8xy$  (c)  $-8xy$  (d)  $-15xy$

(5) If the fraction  $\frac{x-a}{x+3}$  is the multiplicative inverse of  $\frac{x+3}{x+5}$ , then  $a = \dots\dots\dots$

- (a) -5 (b) -3 (c) 3 (d) 5



(٦) If A and B are two mutually exclusive events, then  $P(A \cap B)$  equals .....

(a)  $\emptyset$

(b) zero

(c)  $\frac{1}{2}$

(d) 1

2 Find  $n(X)$  in the simplest form showing the domain of  $n$  where :

(1)  $n(X) = \frac{x^2 + x}{x^2 - 1} - \frac{x - 5}{x^2 - 6x + 5}$

(2)  $n(X) = \frac{x^2 + 2x}{x^3 - 27} \times \frac{x^2 + 3x + 9}{x + 2}$

3 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the following two equations graphically :

$y = x + 4$  ,  $y + x = 4$

[b] By using the formula find in  $\mathbb{R}$  the solution set of the equation :  $2x^2 - 5x - 1 = 0$  approximating the result to the nearest one decimal.

4 [a] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the following two equations :


$x - y = 1$  ,  $x^2 - xy = 0$

[b] Use the opposite Venn diagram and find :

(1)  $P(A \cap B)$

(2)  $P(A \cup B)$

(3)  $P(A - B)$




5 [a] If the domain of the function  $n$  where  $n(X) = \frac{b}{x} + \frac{9}{x+a}$  is  $\mathbb{R} - \{0, 3\}$

,  $n(6) = 7$  find the values of  $a, b$

[b] If  $n_1(X) = \frac{1}{x+1}$  ,  $n_2(X) = \frac{x^2 - x + 1}{x^3 + 1}$  , then prove that :  $n_1 = n_2$

## 23 North Sinai Governorate



Answer the following questions : (Calculators are permitted)

1 Choose the correct answer from those given :

(1) The multiplicative inverse of  $\frac{\sqrt{2}}{3}$  is .....

(a)  $-\frac{\sqrt{2}}{3}$

(b)  $\frac{3\sqrt{2}}{2}$

(c)  $\frac{2\sqrt{3}}{3}$

(d)  $\frac{\sqrt{3}}{2}$

(2) The S.S. of the two equations :  $x - 2y = 1$  ,  $3x + y = 10$  in  $\mathbb{R} \times \mathbb{R}$  is .....

(a)  $\{(5, 2)\}$

(b)  $\{(2, 4)\}$

(c)  $\{(1, 3)\}$

(d)  $\{(3, 1)\}$

(3) Twice its square the number  $\frac{1}{2}$  is .....

- (a)  $-\frac{1}{2}$  (b)  $\frac{1}{2}$  (c)  $\frac{1}{4}$  (d) 1

(4) The domain of the function  $f : f(x) = \frac{x-2}{7}$  is .....

- (a)  $\mathbb{R}$  (b)  $\mathbb{R} - \{2\}$  (c)  $\mathbb{R} - \{7\}$  (d)  $\mathbb{R} - \{2, 7\}$

(5)  $x^2 + kx + 9$  is a perfect square if  $k =$  .....

- (a) 3 (b) -3 (c)  $\pm 3$  (d)  $\pm 6$

(6) If the probability of failure of a student is 0.4 , then the probability of his success is .....

- (a) zero (b) 1 (c)  $\frac{2}{5}$  (d)  $\frac{3}{5}$

**2** [a] Using the general formula , find in  $\mathbb{R}$  the solution set of the equation :

$$x^2 - 2x - 6 = 0$$

[b] Find  $n(x)$  in the simplest form showing the domain of  $n$  where :

$$n(x) = \frac{x}{x-4} - \frac{x+4}{x^2-16}$$

**3** [a] Find in  $\mathbb{R} \times \mathbb{R}$  the S.S. of the following two equations :

$$x - y = 2 \quad , \quad x^2 - 5y = 4$$

[b] If  $n(x) = \frac{x^2 + 3x}{x^2 + x - 6}$

- (1) Find :  $n^{-1}(x)$  and find the domain of  $n^{-1}$  (2) If  $n^{-1}(x) = 2$  , find value of  $x$

**4** [a] Find in  $\mathbb{R} \times \mathbb{R}$  the S.S of the following two equations graphically :

$$y = 2x - 3 \quad , \quad x + 2y = 4$$

[b] Find  $n(x)$  in the simplest form showing the domain of  $n$  where :

$$n(x) = \frac{x^3 - 8}{x^2 - 6x + 5} \div \frac{x^3 + 2x^2 + 4x}{2x^2 + x - 3}$$


**5** [a] A bag contains 15 balls numbered from 1 to 15 , if a ball is drawn randomly , if the event A is getting an odd number and the event B is getting a prime number

Find : (1)  $P(A)$  (2)  $P(B)$  (3)  $P(A - B)$

[b] If  $n_1(x) = \frac{2x}{2x+4}$  ,  $n_2(x) = \frac{x^2 + 2x}{x^2 + 4x + 4}$

Prove that :  $n_1 = n_2$





Answer the following questions : (Calculator is permitted)

1 Choose the correct answer from those given :

(1)  $3^{-2} = \dots\dots\dots$

- (a)  $-9$  (b)  $\frac{-1}{9}$  (c)  $\frac{1}{9}$  (d)  $9$

(2) If A and B are two mutually exclusive events in a random experiment , then  $P(A \cap B) = \dots\dots\dots$

- (a) zero (b)  $\emptyset$  (c)  $1$  (d)  $\{0, 1\}$

(3) The solution set of the inequality :  $x \leq 1$  in  $\mathbb{N}$  is  $\dots\dots\dots$

- (a)  $\{1\}$  (b)  $\{0\}$  (c)  $\{0, 1\}$  (d)  $\{0, 1, -1, \dots\}$

(4) The set of zeroes of  $f$  where  $f(x) = \frac{x^2 - 9}{x - 2}$  is  $\dots\dots\dots$

- (a)  $\{2\}$  (b)  $\mathbb{R} - \{2\}$  (c)  $\{3, -3\}$  (d)  $\{3, -3, 2\}$

(5) If  $n(x) = \frac{x - 7}{x + 3}$ , then the domain of  $n^{-1}$  is  $\dots\dots\dots$

- (a)  $\mathbb{R}$  (b)  $\mathbb{R} - \{-3\}$  (c)  $\mathbb{R} - \{-3, 7\}$  (d)  $\mathbb{R} - \{7\}$

(6) The point of intersection of the two straight lines :  $y = 2$  and  $x + y = 6$  is  $\dots\dots\dots$

- (a)  $(2, 6)$  (b)  $(2, 4)$  (c)  $(4, 2)$  (d)  $(6, 2)$

2 [a] Find the common domain in which the two functions  $f_1$  and  $f_2$  are equal where :

$$f_1(x) = \frac{x^2 + 3x + 2}{x^2 - 4}, \quad f_2(x) = \frac{x^2 - 1}{x^2 - 3x + 2}$$

[b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set to the following two equations graphically :

$$y = x + 4, \quad x + y = 4$$

3 [a] Find  $f(x)$  in the simplest form , showing the domain of  $f$  where :

$$f(x) = \frac{x^2 - x}{x^2 - 1} + \frac{x + 5}{x^2 + 6x + 5}$$

[b] Find in  $\mathbb{R}$  the solution set of the equation :  $x^2 - 2x - 6 = 0$

approximating the result to the nearest two decimals.

- 4 [a] Find  $n(x)$  in the simplest form showing the domain of  $n$  where :

$$n(x) = \frac{x^2 - 3x + 2}{x^2 - 1} \div \frac{3x - 5}{x^2 - 4x - 5}$$

- [b] Find in  $\mathbb{R} \times \mathbb{R}$  the solution set of the two equations :

$$y = x - 3 \quad , \quad x^2 + y^2 = 17$$

- 5 [a] If the set of zeros of the function  $f$  where :

$$f(x) = ax^2 + bx + 8 \text{ is } \{2, 4\} \text{ Find the value of } a \text{ and } b$$

- [b] If  $A$  and  $B$  are two events in a random experiment

$$, P(A) = 0.8 \quad , \quad P(B) = 0.7 \text{ and } P(A \cap B) = 0.6$$

Find : (1) The probability of non occurrence of the event  $A$

(2) The probability of occurrence of at least one of the events.